

REVIEW

Fundamental Mechanisms of Immune Checkpoint Blockade Therapy

Spencer C. Wei¹, Colm R. Duffy¹, and James P. Allison^{1,2}**ABSTRACT**

Immune checkpoint blockade is able to induce durable responses across multiple types of cancer, which has enabled the oncology community to begin to envision potentially curative therapeutic approaches. However, the remarkable responses to immunotherapies are currently limited to a minority of patients and indications, highlighting the need for more effective and novel approaches. Indeed, an extraordinary amount of preclinical and clinical investigation is exploring the therapeutic potential of negative and positive costimulatory molecules. Insights into the underlying biological mechanisms and functions of these molecules have, however, lagged significantly behind. Such understanding will be essential for the rational design of next-generation immunotherapies. Here, we review the current state of our understanding of T-cell costimulatory mechanisms and checkpoint blockade, primarily of CTLA4 and PD-1, and highlight conceptual gaps in knowledge.

Significance: This review provides an overview of immune checkpoint blockade therapy from a basic biology and immunologic perspective for the cancer research community. *Cancer Discov*; 8(9); 1069–86. ©2018 AACR.

INTRODUCTION

Immune checkpoint blockade therapies are now FDA approved for the treatment of a broad range of tumor types (Table 1), with approval likely for additional indications in the near future. The realization of long-term durable responses in a subset of patients represents a transformative event. Since the 2011 FDA approval of ipilimumab (anti-CTLA4) for the treatment of metastatic melanoma, 5 additional checkpoint blockade therapies, all targeting the PD-1/PD-L1 axis, have been approved for the treatment of a broad range of tumor types. Additionally, ipilimumab plus nivolumab (anti-PD-1) combination therapy has been approved for the treatment of advanced melanoma with favorable outcomes compared with either monotherapy. However, as we look to the future and aspire to extend these remarkable responses to more patients and tumor types, many aspects of T-cell activation and the mechanisms of checkpoint blockade remain to be understood. Here, we review how the negative costimulatory molecules CTLA4 and PD-1 attenuate T-cell activation. We also discuss current

dogma and recent conceptual advances related to the mechanisms of action of anti-PD-1 and anti-CTLA4 therapies in the context of antitumor immunity. These discussions highlight the importance of understanding the underlying fundamental biological phenomena for effective translational and clinical research. In the context of the current landscape of cancer immunotherapy, fully understanding how anti-CTLA4 and anti-PD-1 checkpoint blockade therapies work will be critical for effectively combining them with other immunotherapeutic, chemotherapeutic, and targeted approaches.

Immune checkpoint blockade removes inhibitory signals of T-cell activation, which enables tumor-reactive T cells to overcome regulatory mechanisms and mount an effective antitumor response (1–3). Such regulatory mechanisms normally maintain immune responses within a desired physiologic range and protect the host from autoimmunity. Immunologic tolerance is achieved through multiple distinct mechanisms that can be defined as central and peripheral. Central tolerance is mediated through clonal deletion of high-affinity self-reactive clones during negative selection in the thymus. However, because self-reactivity is selected for during positive selection in the thymus, additional mechanisms are required to restrain autoreactivity. Peripheral tolerance is mediated through a variety of mechanisms, including regulatory T cells (Treg), T-cell anergy, cell-extrinsic tolerogenic signals, and peripheral clonal deletion. The immune system exerts a strong selective pressure throughout tumor progression, leading to immune tumor editing (4). As a result, malignant tumors often co-opt immune suppressive and tolerance mechanisms to avoid immune destruction. Immune

¹Department of Immunology, The University of Texas MD Anderson Cancer Center, Houston, Texas. ²Parker Institute for Cancer Immunotherapy, The University of Texas MD Anderson Cancer Center, Houston, Texas.

Corresponding Authors: Spencer C. Wei, The University of Texas MD Anderson Cancer Center, 7455 Fannin Street, Unit 901, Houston, TX 77054. Phone: 713-794-1117; Fax: 713-563-3276; E-mail: scwei@mdanderson.org; and James P. Allison, E-mail: jallison@mdanderson.org

doi: 10.1158/2159-8290.CD-18-0367

©2018 American Association for Cancer Research.

Table 1. Summary of the tumor types for which immune checkpoint blockade therapies are FDA-approved

Tumor type	Therapeutic agent	FDA approval year
Melanoma	Ipilimumab	2011
Melanoma	Nivolumab	2014
Melanoma	Pembrolizumab	2014
Non-small cell lung cancer	Nivolumab	2015
Non-small cell lung cancer	Pembrolizumab	2015
Melanoma (BRAF wild-type)	Ipilimumab + nivolumab	2015
Melanoma (adjuvant)	Ipilimumab	2015
Renal cell carcinoma	Nivolumab	2015
Hodgkin lymphoma	Nivolumab	2016
Urothelial carcinoma	Atezolizumab	2016
Head and neck squamous cell carcinoma	Nivolumab	2016
Head and neck squamous cell carcinoma	Pembrolizumab	2016
Melanoma (any BRAF status)	Ipilimumab + nivolumab	2016
Non-small cell lung cancer	Atezolizumab	2016
Hodgkin lymphoma	Pembrolizumab	2017
Merkel cell carcinoma	Avelumab	2017
Urothelial carcinoma	Avelumab	2017
Urothelial carcinoma	Durvalumab	2017
Urothelial carcinoma	Nivolumab	2017
Urothelial carcinoma	Pembrolizumab	2017
MSI-high or MMR-deficient solid tumors of any histology	Pembrolizumab	2017
MSI-high, MMR-deficient metastatic colorectal cancer	Nivolumab	2017
Pediatric melanoma	Ipilimumab	2017
Hepatocellular carcinoma	Nivolumab	2017
Gastric and gastroesophageal carcinoma	Pembrolizumab	2017
Non-small cell lung cancer	Durvalumab	2018
Renal cell carcinoma	Ipilimumab + nivolumab	2018

NOTE: A summary of the tumor indications, therapeutic agents, and year of FDA approval for immune checkpoint blockade therapies. FDA approval includes regular approval and accelerated approval granted as of May 2018. Ipilimumab is an anti-CTLA4 antibody. Nivolumab and pembrolizumab are anti-PD-1 antibodies. Atezolizumab, avelumab, and durvalumab are anti-PD-L1 antibodies. Tumor type reflects the indications for which treatment has been approved. Only the first FDA approval granted for each broad tissue type or indication for each therapeutic agent is noted. In cases where multiple therapies received approval for the same tumor type in the same year, agents are listed alphabetically.

Abbreviations: MSI, microsatellite instability; MMR, mismatch repair.

checkpoint blockade inhibits T cell–negative costimulation in order to unleash antitumor T-cell responses that recognize tumor antigens. Importantly, the development of immune checkpoint blockade therapies was predicated on basic research that identified key regulatory mechanisms of T-cell activation. However, there remains much to be understood about these mechanisms, further insight into which will be essential for the rational development of immunotherapeutic approaches. The current clinical landscape of cancer immunotherapy and mechanisms of resistance to immunotherapy have been recently reviewed (5–7). Here,

we primarily discuss what is known about the regulatory mechanisms of CTLA4 and PD-1 and the therapeutic implications of these insights.

MECHANISMS OF CTLA4-MEDIATED NEGATIVE COSTIMULATION

CTLA4 expression and function is intrinsically linked with T-cell activation. CTLA4 is immediately upregulated following T-cell receptor (TCR) engagement (signal 1), with its expression peaking 2 to 3 days after activation (8, 9). CTLA4 dampens




Figure 1. Molecular mechanisms of CTLA4 and PD-1 attenuation of T-cell activation. Schematic of the molecular interactions and downstream signaling induced by ligation of CTLA4 and PD-1 by their respective ligands. The possibility of additional downstream cell-intrinsic signaling mechanisms is highlighted for both CTLA4 and PD-1.

TCR signaling through competition with the costimulatory molecule CD28 for the B7 ligands B7-1 (CD80) and B7-2 (CD86), for which CTLA4 has higher avidity and affinity (refs. 10–12; Fig. 1). Because both B7-1 and B7-2 provide positive costimulatory signals through CD28 (refs. 13; signal 2), competitive inhibition of both molecules by CTLA4 is necessary to effectively attenuate T-cell activation. CD28 and CTLA4 also display rapid binding kinetics with B7-1 (12), which, coupled with differences in binding strengths, allows for swift competitive inhibition by CTLA4. In addition to upregulation of CTLA4 expression upon T-cell activation, CTLA4 contained in intracellular vesicles is rapidly trafficked to the immunologic synapse (14). The degree of CTLA4 recruitment to the immunologic synapse correlates directly with TCR signal strength. Once trafficked to the immunologic synapse, CTLA4 is stabilized by B7 ligand binding, allowing it to accumulate and effectively outcompete CD28 (15). Through this mechanism, CTLA4 attenuates positive costimulation by CD28 and thus limits CD28 downstream signaling, which is primarily mediated by PI3K and AKT (16, 17). This results in robust regulation of TCR signal amplitude and, thus, T-cell activity. Because CTLA4-negative costimulation is intrinsically linked to expression of B7 ligands and CD28-mediated positive costimulation, CTLA4 primarily functions to regulate T-cell activity at sites of T-cell priming (e.g., secondary lymphoid organs). In addition to this core function, CTLA4 also attenuates T-cell activation in peripheral tissues given that B7 ligands are constitutively expressed to varying degrees by antigen-presenting cells (APC) but can also be expressed by activated T cells. Because of its central role in regulating T-cell activation, negative costimulation by CTLA4 is critical for tolerance. Reflective of this, biallelic genetic deletion of *Ctla4* leads to massive lymphoproliferation that mice succumb to at 3 to 4 weeks of age (18–20).

In addition to the cell-intrinsic functions through which CTLA4 primarily attenuates T-cell activity, CTLA4 can modulate T-cell activation through several cell-extrinsic mechanisms. Indicative of cell-extrinsic regulatory mechanisms, the presence of CTLA4-competent T cells is sufficient to prevent lethal lymphoproliferation due to genetic deletion of *Ctla4* (21). The majority of cell-extrinsic suppressive function of

CTLA4 is mediated through Tregs (22, 23). Specific loss of CTLA4 in Tregs is sufficient to induce aberrant T-cell activation and give rise to autoimmunity (24, 25). This indicates that Treg-derived CTLA4 is necessary to maintain immunologic tolerance, although it is unlikely that Treg-derived CTLA4 is sufficient to maintain T cell-mediated tolerance. In terms of a potential molecular mechanism, CTLA4 expressed by Treg cells may attenuate T-cell activation in a cell-extrinsic manner by limiting the availability of the B7 ligands B7-1 and B7-2 for CD28-mediated positive costimulation of nearby effector T cells. CTLA4 also has cell-extrinsic contributions within the effector compartment. CTLA4 expressed by effector T cells can compete for B7 ligands in *trans* (26). Additionally, it has been reported that CTLA4 can also act to limit the overall availability of B7 ligands through transendocytosis of B7 ligands from APCs (27). The degree to which these cell-extrinsic processes contribute to T-cell tolerance remains to be fully resolved, particularly in the context of tumor immunity.

Recent work by Sharpe and colleagues demonstrated that genetic loss of CTLA4 in Tregs in adulthood surprisingly confers resistance to experimental autoimmune encephalomyelitis (EAE; ref. 28). Conditional deletion of *Ctla4* in Tregs is necessary and sufficient to confer resistance to EAE, suggesting that unrestrained peripheral Treg expansion and/or increased Treg activation can prevent autoimmunity. A significant implication of this finding is that Treg depletion may counter expansion of Treg cells induced by CTLA4 blockade and thus lead to enhanced efficacy of anti-CTLA4 therapy. This observation also raises the possibility that CTLA4 has differential functions in conventional and regulatory T cells during development and in adulthood. Alternatively, the apparently discordant observations between global and conditional *Ctla4* knockout mice may be the result of the intrinsic difference in antigen affinity of conventional T cells and Tregs. Tregs are selected for higher-affinity TCR for self-peptide MHC complexes, and thus because CTLA4 expression correlates with TCR signal strength, have concomitantly higher CTLA4 (29, 30). This mechanism attenuates strong TCR signals, allowing medium-strength TCR signals to also result in robust T-cell activation. As a result, loss of CTLA4 may disproportionately affect T cells with high-affinity antigen receptors. These findings may support the strength of signal model proposed more than 20 years ago (31). Moving forward, it will be critical to precisely dissect the function of downstream signaling components and assess their relative functional contribution to CTLA4-mediated regulation of T-cell activity.

MECHANISMS OF PD-1-MEDIATED ATTENUATION OF T-CELL ACTIVITY

The primary biological functions of PD-1 are to maintain peripheral tolerance and to maintain T-cell responses within a desired physiologic range. Because the PD-1/PD-L1 regulatory system is induced by immune responses (discussed in greater detail below), this forms a negative feedback loop to attenuate local T-cell responses and minimize tissue damage. PD-1 regulates T-cell activation through interaction with PD-L1 and PD-L2 (refs. 32–34; Fig. 1). PD-1 is expressed upon activation of T and B lymphocytes (35). Because of

the expression of its ligands, which are widely expressed in nonlymphoid tissues, PD-1 acts primarily to dampen T-cell activation in the periphery (36). PD-L1 expression, and to a lesser degree expression of PD-L2, is induced in response to inflammatory cytokines such as IFN γ (32, 33). Thus, PD-1 regulation of T-cell activity occurs in response to cytolytic and effector T-cell function [e.g., CD8 cytotoxic T lymphocyte and type 1 helper (Th1) CD4 T cells] in an inducible manner. Upon engagement with PD-L1 and PD-L2, PD-1 is thought to primarily transmit a negative costimulatory signal through the tyrosine phosphatase SHP2 to attenuate T-cell activation. The recruitment of SHP2 directly attenuates TCR signaling via dephosphorylation of proximal signaling elements (37). This molecular mechanism reflects a dichotomy in modes of regulation utilized by CTLA4 and PD-1 engagement (38). These data indicate that in contrast to CTLA4-mediated regulation, PD-1 directly regulates TCR signaling to attenuate T-cell activity. However, recent evidence indicates that CD28 is a primary target for PD-1-induced attenuation of T-cell signaling (39). These studies utilized a cell-free membrane reconstitution model to examine functional relationships during T-cell activation and reveal that PD-1 leads to preferential dephosphorylation of CD28 rather than the TCR, via recruitment of SHP2. This suggests that both CTLA4 and PD-1, at least in part, act through a similar molecular mechanism of attenuating CD28-mediated costimulation (Signal 2). Thus, modulation of CD28 signaling could represent a functional convergence point of CTLA4- and PD-1-mediated regulation. Interestingly, recent findings indicate that SHP2 is not essential for responses to anti-PD-1 therapy or induction of T-cell exhaustion *in vivo* (40). This is suggestive of functional redundancy in the signaling pathways downstream of PD-1. Such redundancy is most likely mediated through redundant phosphatases (e.g., SHP1) but alternatively could be mediated through wholly distinct mechanisms. It is critical to further define the immediate signaling events downstream of CTLA4 and PD-1 to distinguish shared and distinct molecular mechanisms of these T-cell regulatory pathways.

Functionally, PD-1 is essential for homeostatic maintenance of peripheral tolerance as evidenced by the autoimmune pathologies that arise upon genetic deletion of *Pdcd1* (encoding PD-1). As an example, genetic loss of *Pdcd1* leads to development of lupus-like autoimmune pathology in aged C57BL/6 mice and autoimmune dilated cardiomyopathy in BALB/c mice (41, 42). The strong murine strain dependency of the PD-1 knockout phenotype raises the possibility that the observed autoimmunity may be driven by recognition of strain-specific antigens in the absence of PD-1 inhibitory signaling; however, this remains to be definitively tested. It is a critical point that although PD-1 is often used as a marker of exhaustion, it is not sufficient to define a functionally exhausted population. PD-1 is a marker of activated T cells, of which exhausted T cells are a subset. Exhausted T cells are often defined by coexpression of PD-1, LAG3, and TIM3. However, an essential distinction is that exhausted T cells (phenotypically defined) are still functionally active, but harbor reduced capacity. Thus, for example, exhausted CD8 T cells are still able to contribute to antitumor immune responses, but are likely less potent on a per-cell basis.

T-cell exhaustion is an important mechanism that limits T-cell activity in the presence of chronic antigen stimulation and acts to preserve T-cell clones that would otherwise perish under such conditions due to activation-induced cell death. Consistent with this notion, persistent PD-1 signaling induces metabolic restriction, which is a functional driver of T-cell exhaustion (43). Upon ligation, PD-1 attenuates glycolysis but simultaneously promotes fatty-acid oxidation and lipid catabolism, thus inducing a switch in energy derivation (44). In contrast, ligation of CTLA4 attenuates glycolysis independent of regulation of lipid metabolism. Interestingly, this metabolic switch is involved in determining T-cell effector versus memory fate and is driven in part by mitochondrial regulation (45). Such changes are likely driven by the changes in gene expression and epigenetic regulation that are induced by continuous PD-1 engagement. Indeed, chronic antigen stimulation in viral systems leads to dramatic changes in gene regulation and stable epigenetic reprogramming of T cells (46, 47). Together, such transcriptional, epigenetic, and metabolic changes define the exhausted T-cell state. Recent evidence suggests that such epigenetic changes can prevent the rescue of an exhausted state by checkpoint blockade and attenuate tumor responses to therapy (48).

Emerging evidence has also identified new functional roles for the PD-1/PD-L1 signaling axis. For example, macrophage expression of PD-L1 may lead to active eviction of T cells from the tumor microenvironment (49). This suggests that in addition to regulation of T-cell activation and cytolytic capacity, PD-1 signaling may also regulate T-cell trafficking and migration. Furthermore, it has been reported that PD-1 may also have tumor cell-intrinsic function (50). Future studies are needed to determine the degree to which such “noncanonical” mechanisms contribute to therapeutic efficacy.

MECHANISMS OF NEGATIVE COSTIMULATION VERSUS MECHANISMS OF CHECKPOINT BLOCKADE

Insights into the normal biological roles and molecular mechanisms of costimulatory molecules undoubtedly inform our understanding of mechanisms of action of cancer therapeutics targeting these molecules. Differences will remain however, given the properties of specific therapeutics (e.g., antibody isotype, off-target recognition, kinetics) as well as the property of cancer being self-derived but ideally recognized as foreign by the immune system. Based on our understanding of how the molecules themselves act to attenuate T-cell activity, it is thought that anti-CTLA4 and anti-PD-1 primarily act at different stages of the cancer-immunity cycle (51). Conceptually, the current model posits that CTLA4 blockade primarily acts at sites of priming in which CD28-positive costimulation is involved (e.g., tumor-draining lymph nodes) whereas PD-1 blockade primarily acts in inflamed peripheral tissues (e.g., tumor; Fig. 2). Recent evidence, discussed below, raises the possibility that the mechanisms of action of CTLA4 and PD-1 blockade are not limited to only these tissue sites.

Many of the principles and lessons learned in viral systems are applicable to tumor immunity, as cancer is highly




Figure 2. Schematic of the molecular mechanisms of action of CTLA4 and PD-1 blockade. The step-wise progression of T-cell activation, attenuation by normal regulatory mechanisms, and release of such negative regulation by therapeutic intervention using anti-CTLA4 or anti-PD-1 antibodies is outlined (left). In addition to cell-intrinsic enhancement of effector function, several additional mechanisms are thought to contribute to the efficacy of anti-CTLA4 and anti-PD-1 therapy (right). These include antibody-mediated depletion of Tregs, enhancement of T-cell positive costimulation within the tumor microenvironment, blockade of host-derived PD-L1 signals from nontumor cells in the microenvironment (as opposed to tumor cell-derived PD-L1), and blockade of interactions between PD-L1 and B7-1.

analogous to infectious disease contexts in which chronic antigen stimulation results in T-cell exhaustion (52). For example, blockade of PD-1 is sufficient to enhance the activity of exhausted T cells in the context of chronic viral infection, leading to viral clearance (53). Recent findings demonstrate that CD28 costimulation is necessary for responses to PD-1 blockade in the settings of both viral infection and tumor rejection (54). Together, these findings indicate that additional positive costimulation is required for therapeutic efficacy despite prior activation. This raises the possibility that PD-1 blockade acts not only in peripheral tissues (e.g., tumor) but also in sites of priming. The mechanisms of action of PD-1 and CTLA4 blockade and of the normal biological functions of these molecules are highly complex and clearly not fully understood. It is likely that subtle nuances in pertinent aspects of such mechanisms (e.g., timing, kinetics, target cell type, cognate antigen availability, anatomic location) will have profound impact on the final biological outcomes.

MECHANISMS OF ACTION OF CTLA4 BLOCKADE-INDUCED TUMOR REJECTION

CTLA4 blockade is thought to induce tumor rejection through a number of distinct mechanisms. The primary mechanism seems to be through direct blockade of CTLA4 competition for B7-1 and B7-2 costimulatory ligands, which allows for unrestrained CD28-mediated positive costimulation. Indeed, crystallographic structural analyses of the ipilimumab:CTLA4 complex reveal that the ipilimumab binding epitope overlaps with the B7 interaction domain, indicating that steric inhibition of B7 interactions underlies the primary mechanism of

action of ipilimumab (55). Because tumor cells do not express B7 ligands, this action largely occurs in tumor-draining lymph nodes in which tumor antigens can be cross-presented by APCs to prime tumor-reactive T cells. It is also feasible that APCs within the tumor microenvironment may also cross-present tumor antigens to activate cognate tumor-reactive T cells. In either case, tumor cell death is required to release tumor cell antigens (e.g., neoantigens, tumor-associated antigens) that can be subsequently processed and presented by APCs. In the context of effective antigen presentation, CTLA4 blockade then enhances CD28 costimulation and thus activation. The extent to which APCs can directly process and cross-present tumor antigens within the tumor microenvironment to prime (or reprime) T cells *in situ* remains unclear. An interesting possibility is that antigen presentation may be taking place in tumor-associated tertiary lymphoid structures (TLS), the presence of which is generally associated with improved survival (56, 57). The role of TLS in antitumor immunity is complex and likely context-dependent however, as Treg populations within TLS have been shown to suppress antitumor T-cell responses (58). Understanding when and where antitumor T cells are primed and subsequently regulated (and thus potentially sensitive to checkpoint blockade therapy) remains a critical open question.

Emerging evidence suggests that anti-CTLA4 does not impose a generalized effect on all T cells. CTLA4 blockade leads to specific expansion of tumor neoantigen-specific CD8 T cells within the tumor microenvironment, but not secondary lymphoid organs (59). Consistent with this notion, anti-CTLA4 leads to expansion of specific tumor-infiltrating T-cell populations including a subset of phenotypically exhausted CD8 T cells and a PD-1⁺ICOS⁺TBET⁺ Th1-like CD4 effector

T-cell population (60). This population appears to differ from canonical Th1 cells due to coexpression of ICOS and PD-1, which are markers of T follicular helper cells. Whether such cells reflect a distinct type of T-cell that emerges after therapy or, alternatively, an activated phenotype of a preexisting minor population remains to be fully determined. These findings are supported by clinical observations of expansion of ICOS⁺ CD4 effector T cells following ipilimumab therapy in multiple tumor types (61–64) as well as following treatment with tremelimumab, another anti-CTLA4 antibody (65). Thus, the expansion of ICOS⁺ CD4 effector T cells may be used as a pharmacodynamic marker of anti-CTLA4 therapy (66). Furthermore, the expansion of specific types of CD4 effector T cells raises the possibility that anti-CTLA4 not only enhances T-cell activation, but may also affect T-cell differentiation. The extent to which effects on T-cell lineage choices may contribute to the mechanisms and efficacy of immune checkpoint blockade therapies remains unclear. Notwithstanding, together these and other data discussed below indicate that CTLA4 blockade enhances antitumor immunity through modulation and expansion of particular T-cell populations.

In addition to these mechanisms of CTLA4 blockade-induced tumor rejection, depletion of Treg populations has also been identified as a mechanism of action of anti-CTLA4 therapy in murine tumor models (67–69). Treg depletion contributes partially to antitumor efficacy, as significant therapeutic benefit was still observed in Fc-gamma-RIV knockout C57BL/6 hosts (67). Interestingly, Treg depletion appears to be carried out with differential efficiency depending on the context. Treatment with depleting clones of anti-CTLA4 decreases intratumoral but not peripheral populations of Treg cells (68). This can be explained by the increased expression of CTLA4 by intratumoral Tregs, or alternatively by differences in the abundance and activity of Fc receptor-expressing cell populations in each context (e.g., tumor-associated macrophages). A recent report asserts that the efficacy of anti-CTLA4 is completely independent of its regulation of B7 ligand interactions and, instead, derives solely from antibody-mediated Treg depletion (70). This is a conclusion that is in conflict with numerous lines of prior evidence, and the experimental observations presented therein are insufficient to definitively arrive at this conclusion. For example, indirect readouts are used to infer molecular interactions and arrive at key observations such as the dispensability of B7 ligands for the therapeutic efficacy of anti-CTLA4. These conflicting findings may be explained by technical limitations of the biological systems utilized rather than biological independence of anti-CTLA4 and inhibition of B7 ligand interactions. Furthermore, this finding is in direct conflict with structural biology studies that revealed that ipilimumab binds CTLA4 precisely in the B7 interaction domain to mediate steric hindrance (55). Although other prior studies (discussed below in greater detail) do suggest that Treg depletion contributes to the mechanism of action of anti-CTLA4, a large body of work strongly implicates regulation of B7 ligand interactions as a critical mechanism. Notably, blockade of both effector and regulatory T-cell compartment-derived CTLA4 is required for effective tumor rejection (71). Notwithstanding, it is clear that detailed and careful determination of the relative contri-

bution of each of the mechanisms of action of anti-CTLA4 therapies, particularly in the context of human immunity, is required.

The relative contribution of cell-intrinsic enhancement of effector function versus Treg depletion to the efficacy of ipilimumab in humans remains unclear. Ipilimumab was purposefully selected to be a blocking antibody based on the understanding that loss of CTLA4 would lead to enhanced T-cell activity. Thus, although ipilimumab is a fully human IgG1 antibody, it was not developed to be a depleting antibody. Consistent with this, there is not definitive evidence of Treg depletion in patients treated with ipilimumab. It is difficult to perform the studies required to resolve this issue of contention, as such studies are hindered by the necessity for paired pre- and post-therapy sampling, high tumor heterogeneity (in terms of immune infiltration), lack of clarity of how to best normalize quantification, and because FOXP3 is expressed in activated effector T cells in humans. It has been reported that ipilimumab can induce antibody-dependent cellular cytotoxicity (ADCC)-mediated killing of Tregs by nonclassic monocytes in *ex vivo* cultures (72). Moreover, recent evidence indicates that germline presence of a high-affinity polymorphism in the Fc receptor (CD16a-V158F) is associated with improved responses to ipilimumab (73). This suggests that Fc-mediated cell depletion functionally contributes in part to the mechanism of ipilimumab. In contrast, the similarity in response rates of two anti-CTLA4 antibodies (tremelimumab and ipilimumab) despite different antibody isotypes supports the notion that efficacy derives from enhancement of effector function rather than depletion. Tremelimumab is a fully human IgG2 antibody, whereas ipilimumab is a fully human IgG1 antibody, which is notable because IgG1 antibodies more effectively mediate ADCC than IgG2 antibodies based on their respective binding affinity for human Fc receptors (74). Although tremelimumab did not reach statistical significance in overall survival at the planned second interim analysis in the phase III clinical trial in metastatic melanoma, follow-up analyses suggest that responses to tremelimumab are roughly comparable to those of ipilimumab (75). Pooled analyses of the phase I and II clinical trials of tremelimumab reveal a 5-year survival rate of 20% (76), which is similar to the 21% 3-year survival rate observed in patients with metastatic melanoma treated with ipilimumab (77). These data support a model in which anti-CTLA4 both enhances cell-intrinsic effector function through blockade and induces Fc-mediated cellular depletion.

Modulation of the TCR repertoire may also contribute to the therapeutic effects of CTLA4 blockade. For example, ipilimumab treatment leads to a remodeling and broadening of the peripheral TCR repertoire (78, 79). Consistent with these findings, ipilimumab therapy broadens the functional reactivity of peripheral blood CD8 T cells for melanoma antigens (80). Interestingly, TCR repertoire broadening is also associated with immune-related adverse events (irAE) due to ipilimumab treatment (81), although it remains to be determined whether the underlying mechanisms and cognate antigens involved in therapeutic efficacy and irAEs are similar. Together, these observations suggest that TCR repertoire broadening due to blockade of CTLA4 has significant clinical relevance. Mechanistically, loss of CTLA4 may lower

the threshold for TCR ligation required for effective T-cell activation given that CTLA4 normally acts to attenuate TCR signal strength. Upon blockade of CTLA4, antigens with low signal strength that are not normally sufficient to generate an effective T-cell response may be allowed to emerge. Such T-cell clones could recognize tumor-specific antigens (e.g., subdominant neoantigens) or tumor-associated antigens. In addition, the activity of high-affinity tumor-reactive clones would also be boosted by blockade of CTLA4 through this mechanism. Taken together, our understanding of the biology underlying CTLA4 indicates that its blockade acts to increase T-cell costimulation in multiple distinct ways, resulting in more robust activation of tumor-reactive T cells. Multiple lines of evidence indicate that tumor mutational burden (TMB) is associated with improved responses to checkpoint blockade (82–86). This supports a model in which neoantigens are a major driver of tumor immunogenicity. On the other hand, some tumor types such as renal cell carcinoma exhibit responsiveness to checkpoint blockade despite low mutational burden (87). It remains a possibility that low-TMB tumors that respond to checkpoint blockade harbor low numbers of highly immunogenic tumor-specific neoantigens. Alternatively, mechanistically distinct types of antitumor immune responses may underlie responses of low-TMB tumor types. Relatedly, the relative contribution of public antigens (shared; e.g., overexpressed genes) and private antigens (tumor-specific; e.g., neoantigens) to antitumor immune responses remains a key outstanding question.

MECHANISMS OF ACTION OF PD-1 BLOCKADE-INDUCED TUMOR REJECTION

PD-1 blockade is able to induce tumor rejection through reinvigoration of CD8 T cells, leading to both increased functional activity and frequency. Blockade of the PD-1 signaling axis prevents PD-1-mediated attenuation of proximal TCR signaling, allowing for restoration of activity of exhausted CD8 effectors. Thus, despite continued PD-L1 expression within the tumor microenvironment, exhausted T cells are able to be reinvigorated and mount an effective immune response. Clinical evidence supports a model in which blockade of the PD-1 signaling axis is most effective in tumors in which an endogenous T-cell response has already been elicited but is suppressed through PD-1 engagement by its ligands PD-L1 and PD-L2 (88, 89). However, the response of some PD-L1-negative tumors indicates that the presence of a preexisting immune response, as arbitrarily defined by the presence of tumor-infiltrating T cells, is not required for tumor rejection to be induced by PD-1 blockade. Recent evidence suggests that a subset of CXCR5⁺ PD-1⁺ CD8 T cells is responsible for immediate proliferative expansion following PD-1 blockade (90). Longitudinal profiling of peripheral blood from patients treated with anti-PD-1 therapies reveals expansion of PD-1⁺ CD8 T cells with kinetics consistent with this notion (91). The antigen specificity of the T cells that mediate responses to checkpoint blockade therapy remains ill-defined. Recent evidence from a neoadjuvant trial of nivolumab in the context of non-small cell lung cancer supports the notion that anti-PD-1 therapy enhances neoantigen-specific T-cell responses (92). It is likely that only specific T-cell populations (defined

by antigen specificity and/or phenotype) functionally mediate responses to checkpoint blockade therapy. Consistent with this notion, exhausted T cells display a distinct epigenetic profile, and this epigenetic reprogramming can limit T-cell reinvigoration (47, 48, 93, 94). These data suggest that PD-1 blockade may not be sufficient to functionally restore T cells once they reach a threshold level of exhaustion. Recent work reveals a high degree of phenotypic and functional heterogeneity within exhausted CD8 T cells (95). It will be conceptually critical to understand how functional heterogeneity of exhausted T cells affects the mechanisms of action and efficacy of specific checkpoint blockade therapies.

Despite much active investigation and interest in the field, the precise molecular and cellular events that mediate enhancement of antitumor immunity by PD-1 blockade remain not fully understood. Recent studies have revealed subtleties that are likely to have very important consequences for therapeutic efficacy and rational design of new strategies. For example, although PD-1 blockade primarily leads to the expansion of CD8 T cells, CD4 T cells are required for effective responses (96). Although this is not entirely surprising given the critical roles CD4 help plays during a wide range of processes including memory formation and antibody production, this emphasizes the complexity in defining mechanisms of action. In particular, this highlights the distinction between cellular processes that are modulated by therapy and cellular processes that are required for therapeutic efficacy.

It remains unclear what specific aspects of CD4 help are functionally required for clinical responses to checkpoint blockade. In addition to facilitating T-cell memory formation, it is tempting to speculate that CD4 helper T cells may also enhance antitumor immunity by increasing CD8 T-cell and antibody entry into peripheral tissue sites, as has been analogously observed in viral contexts (97, 98). In addition to ambiguities at the cellular level, emerging evidence has shed new insights into molecular mechanisms of PD-1 blockade. In addition to restoring T-cell activity through modulation of TCR signaling and gene expression, blockade of the PD-1 signaling axis is able to reverse the associated metabolic reprogramming to an extent, which in part mediates T-cell reinvigoration (43). Supportive of this finding, anti-PD-1 treatment has been shown to regulate metabolic function based on gene set enrichment analysis of tumor antigen-specific tumor-infiltrating lymphocytes (99). In contrast, CTLA4 blockade primarily leads to changes in genes associated with proliferation and cell cycle. In addition to preventing attenuation of T-cell activation, PD-1 blockade may also act through additional mechanisms that contribute partially to its therapeutic efficacy. For example, it has been reported that tumor cell-intrinsic PD-1 can promote melanoma growth (50).

In addition to direct blockade of PD-1, antibodies targeting PD-L1 are also sufficient to induce immune tumor rejection. Blockade of PD-L1 is thought to largely phenocopy the effect of PD-1 blockade given the dominance in expression of PD-L1. PD-L1 is induced by Th1 cytokines (e.g., IFN γ) whereas PD-L2 is induced by Th2 cytokines (100). This differential regulation may in part explain the efficacy of PD-L1 blockade because Th1-skewed responses would be more favorable for antitumor immune responses. In contrast to anti-PD-1 antibodies, blockade of PD-L1 may also derive part of its efficacy from ADCC. It was recently

demonstrated that Fc receptor binding is important for the efficacy of anti-PD-L1, but not anti-PD-1, antibody therapy-induced tumor regression in murine tumor models (101). Another complicating aspect of the underlying biology is that in addition to canonical binding relationships described, B7-1 and PD-L1 also interact, leading to inhibition of T-cell activity (102). Significantly, these data suggest that anti-PD-1 and anti-PD-L1 therapies are not completely mechanistically equivalent. Recent evidence indicates that host-derived PD-L1 expression is required for the PD-L1 blockade-induced tumor rejection (103, 104). Other evidence indicates, however, that tumor-derived PD-L1 is sufficient to inhibit antitumor immunity via attenuation of CD8 T-cell cytotoxicity (105). How these apparently disparate findings can be integrated remains to be fully understood. Nonetheless, they raise the possibility that PD-L1 can inhibit T cell-mediated tumor cell killing through both cell-autonomous and nonautonomous mechanisms.

THERAPEUTIC COMBINATIONS

Despite the remarkable progress that has been achieved with monotherapies, there is a tremendous need to improve efficacy across tumor types. Understanding which aspects of the tumor microenvironment functionally limit responses to immune checkpoint blockade therapy is an active area of investigation. A primary mechanism is compensatory upregulation of additional immune checkpoint molecules, which limit the therapeutic efficacy of monotherapy approaches. For example, increased expression of PD-L1 (and engagement of PD-1) may in part explain why anti-CTLA4 monotherapy has not resulted in significantly enhanced response rates in tumor types other than melanoma. Consistent with this notion and our understanding that PD-1 and CTLA4 attenuate T-cell activation through distinct mechanisms, combination blockade of PD-1 and CTLA4 improves therapeutic efficacy compared with either monotherapy (106–108). These findings reflect the enhanced efficacy also observed in preclinical models (109). Notably, combination treatment is able to produce complete responses in 36% of patients, and over half of patients with melanoma achieve objective responses (107). Pooled analysis of the 3-year follow-up from the phase II and III clinical trials in melanoma reports a 57% 3-year overall survival in the ipilimumab plus nivolumab group (110). Based on the assumption that durability of responses to combination therapy reaching 3 years will at least equal or exceed that observed in response to ipilimumab monotherapy (77), it is conceivable that greater than half of patients with metastatic melanoma treated with the combination of ipilimumab and nivolumab may achieve long-term responses lasting 10 years or more. Nivolumab plus ipilimumab also improves overall survival versus standard-of-care sunitinib in advanced renal cell carcinoma (111), suggesting that combination therapy may have broad therapeutic efficacy.

Mechanistically, it remains unclear whether the enhanced efficacy of combination anti-PD-1 and anti-CTLA4 therapy is mediated by additive engagement of the cellular and molecular mechanisms of the respective monotherapies or, alternatively, through mechanisms distinct from the component monotherapies. Profiling of peripheral blood supports a model in which PD-1 and CTLA4 act through independ-

ent mechanisms, with combination inhibition of PD-1 and CTLA4 leading to distinct immune responses (112). Similar analyses have interestingly revealed immunologic changes in peripheral B cells associated with the development of immune-related adverse events (113). These observations provide additional support to the notion that combination therapy induces distinct cellular and molecular changes and highlight that these mechanisms may be either direct or indirect. Given that PD-1 and CTLA4 attenuate T-cell activity through separate molecular mechanisms and that blockade of these respective molecules regulate distinct cell populations (60), multiple possible mechanisms may underlie the enhanced efficacy of anti-CTLA4 and anti-PD-1 combination therapy (Fig. 3). Resolving this ambiguity and determining the precise cellular and molecular mechanisms of combination anti-CTLA4 plus anti-PD-1 therapy is critical.

The molecular and cellular mechanisms of anti-CTLA4 and anti-PD-1 monotherapies may provide insights, although it remains unclear the degree to which mechanisms of combination therapy directly reflect those of monotherapies. Given that both CTLA4 and PD-1 have cell-intrinsic regulatory activity, simultaneous blockade of both molecules may lead to functional convergence through enhancement of T-cell activity (whether by coregulation of CD28 or other signaling pathways involved in T-cell activation). Conceptually, this convergence may occur in non-mutually exclusive scenarios (Fig. 3). In the first scenario, CTLA4 and PD-1 are simultaneously targeted on the same cell, leading to an additive increase in CD28 costimulation and T-cell activity. In the second scenario, combination therapy targets T cells at different times with respect to activation and/or trafficking. This model is supported by the distinct kinetics of PD-1 and CTLA4 expression during T-cell activation. Through such a mechanism, combination therapy may broaden the duration and integrated strength of T-cell costimulation by CD28. Both of these scenarios are in part predicated on the assumption that anti-PD-1 and anti-CTLA4 target the same cell population. It remains conceptually unclear whether immune checkpoint blockade reestablishes positive costimulatory levels to normal maximal levels (Fig. 4A), expands the range of T-cell clones able to activate by lowering the costimulatory threshold (Fig. 4B), or, alternatively, whether checkpoint blockade is able to increase activity on a per-cell basis by enhancing costimulatory signals beyond normal physiologic levels (Fig. 4C). A fascinating possibility is that positive costimulation beyond physiologic levels may allow for the acquisition of enhanced cytolytic capabilities or novel properties not displayed by canonical T-cell populations.

In addition to dual engagement of convergent molecular pathways, engagement of distinct cellular biology by anti-CTLA4 and anti-PD-1 may also contribute to the enhanced efficacy of combination therapy. Indeed, anti-CTLA4 but not anti-PD-1 checkpoint blockade leads to the expansion of a tumor-infiltrating ICOS⁺ Th1-like CD4 effector population (60). Consistent with additional reports (90, 91, 114, 115), this suggests that anti-PD-1 primarily acts through targeting of CD8 T-cell populations. This significant difference in the mechanisms of anti-CTLA4 and anti-PD-1 raises the possibility that the enhanced efficacy of combination therapy is due to engagement of multiple distinct populations. Thus,




Figure 3. Potential cellular mechanisms that mediate tumor rejection in response to combination anti-CTLA4 and anti-PD-1 checkpoint blockade. Multiple non-mutually exclusive models of the cellular mechanisms underlying combination anti-CTLA4 plus anti-PD-1 therapy of action are proposed. Models described from left to right: (i) the same T cells may be targeted at the site of priming, leading to enhanced penetrance of effective blockade (i.e., a greater proportion of target cells receive sufficient signal to increase activation) and/or enhanced costimulatory signals beyond normal limits, (ii) different T-cell populations are targeted within the site of priming, potentially leading to synergistic effects through cell-extrinsic processes (e.g., providing CD4 help to CD8 effector T cells), (iii) the same T cells are targeted but with different spatiotemporal kinetics leading to perhaps prolonged costimulatory signaling, and (iv) different T-cell populations are targeted in different tissues (e.g., PD-1 blockade primarily acting on preexisting tumor-infiltrating CD8 T cells whereas CTLA4 acts on CD4 effector T cells in secondary lymphoid organs). T-cell subsets are denoted as “A” and “B” given that the precise populations that are directly targeted remain to be fully defined, particularly in the context of kinetics of therapy and different tissue sites. Conceptually, T-cell “A” and “B” could, for example, represent particular subsets of tumor-specific CD8 T cells and CD4 effector T cells, respectively. Potential effects are noted below each scenario; however, there is certain to be additional aggregate effects and differences between these models. Only secondary lymphoid organs (e.g., draining lymph node) and tumor are described, but other tissue sites may have functional contributions to this process as well.

multiple, non-mutually exclusive mechanisms may facilitate the enhanced efficacy of combination therapy (Fig. 3). Thus, combination therapy may utilize cellular and molecular mechanisms as yet unidentified, which are completely distinct from those that mediate monotherapy-induced tumor rejection. Distinguishing these possibilities will be significant in guiding whether insights into monotherapy mechanisms can be extrapolated to understand how respective combinations work.

The engagement of the CD4 effector compartment resulting in expansion of Th1-like CD4 effectors following anti-CTLA4 but not anti-PD-1 also provides some mechanistic rationale for the possibility that sequential treatment of anti-CTLA4 followed by anti-PD-1 therapy may be advantageous. An increase in CD4 help during the priming and early activation stage as a result of CTLA4 blockade would likely improve T-cell memory development as well as infiltration into peripheral tissues (e.g., tumor). However, upon entry into the tumor microenvironment, Th1 and CD8 effector T cells will induce PD-L1 expression on tumor cells and stromal cells, attenuating T-cell activity. Thus, sequential combination of first CTLA4 and then PD-1 blockade could potentially induce T-cell infiltration of immunologically barren tumors and allow them to maintain effective cytolytic activity within the tumor microenvironment. In contrast, results from a phase II open-label study suggest

that nivolumab followed by ipilimumab has improved efficacy compared with the ipilimumab followed by nivolumab in advanced melanoma (116). These observations remain to be further validated, but nonetheless raise the question of whether sequential therapies can be designed based only on mechanisms of action of monotherapies and also the extent to which properties such as kinetics of response need to be considered. In addition, these observations raise the possibility that anti-CTLA4-induced CD4 help may not be required for rejection of already well-infiltrated tumors (e.g., melanoma). Nonetheless, sequential treatment may minimize the increase in adverse events that are associated with simultaneous combination treatment. On the other hand, simultaneous combination therapy has enhanced overall response rates in melanoma compared with monotherapies (107, 110), and a key unanswered question is whether sequenced therapy has similarly enhanced response rates and long-term efficacy compared with simultaneous combination therapy.

More broadly, the relative contribution of each of the several known molecular mechanisms of CTLA4 and PD-1 blockade to therapeutic efficacy remains unclear. Such differences may manifest in distinct requirements for the induction of effective immune responses in the context of each therapy. For example, several lines of evidence indicate that cross-priming mediated




Figure 4. Potential models of how immune checkpoint blockade restores positive costimulation and modulates T-cell activity. Three non-mutually exclusive theoretical models of how checkpoint blockade regulates the strength of positive costimulation and enhances antitumor immunity. Distinguishing these models will have significant implications for the types and specificities of T cells that are functionally essential for therapeutic efficacy. **A**, In the first model, checkpoint blockade restores the positive costimulatory signaling to levels similar to those reached prior to attenuation (e.g., by PD-1 or CTLA4). Presumably, in this model, enhanced efficacy due to checkpoint blockade would be primarily derived from an increase in the number of activated and cytolytic T cells. **B**, In the second model, increased positive costimulation due to blockade of negative costimulatory molecules lowers the effective threshold required for TCR signal strength. This in effect would allow for activation and expansion of weaker T-cell clones (i.e., low-affinity, low-avidity), which are normally restrained. **C**, In the third model, checkpoint blockade leads to an increase in positive costimulatory signals beyond levels that are achieved in normal scenarios. In this model, enhanced efficacy could be derived from increased number of activated T cells and/or acquisition of new or enhanced functional properties due to super physiologic levels of costimulatory signaling.

by CD103⁺ BATF3-dependent dendritic cells is required for effective antitumor immunity and responses to checkpoint blockade (117–119). Indeed, CD103⁺ dendritic cell populations appear to be the primary cell population to efficiently take up tumor antigen and present it within the draining lymph node (120). Whether the same modes of antigen presentation are important for anti-CTLA4 and anti-PD-1 therapies is unclear.

Mechanistic differences between these therapies may impose distinct requirements in terms of the cellular context and temporal dynamics of antigen presentation.

How immune checkpoint blockade mechanistically interacts with conventional therapies (e.g., surgery, chemotherapy, radiation, targeted therapies) and other immune-based therapies (e.g., chimeric antigen receptor T-cell therapy, other adoptive transfer

approaches, cytokine therapy, personalized tumor vaccines) is a topic of clear relevance and active investigation. For example, radiation treatment and blockade of the PD-1/PD-L1 axis have been shown to have additive effects through nonredundant mechanisms (121, 122). Notably, abscopal responses have been observed following concurrent radiation and CTLA4 blockade, highlighting a potential mechanistic basis for synergistic efficacy (123, 124). Targeted inhibition of immunosuppressive myeloid populations in combination with immune checkpoint blockade therapy leads to enhanced efficacy (125). Additional clinical variables and patient characteristics may also prove to be significant modulators of response to immunotherapy. For example, recent work has elucidated a role for the gut microbiome in defining tumor responses to immunotherapies. Colonization by specific commensal bacteria strains modulates the efficacy of immune checkpoint blockade therapy in preclinical and clinical settings (126–129). This highlights how a diverse set of host properties, in addition to tumor characteristics, can contribute to sensitivity to immunotherapy.

BEYOND CTLA4 AND PD-1

T-cell costimulatory molecules as a functional category represent a large number of proteins belonging to multiple structurally defined superfamilies. The therapeutic potential of many of these targets is now being investigated preclinically and clinically. Among these molecules are LAG3, TIM3, TIGIT, VISTA, and ICOS from the immunoglobulin superfamily (IgSF) and OX40, GITR, 4-1BB, CD40, and CD27 from the tumor necrosis factor receptor superfamily (TNFRSF). However, our collective understanding of the fundamental biological roles of these molecules remains unsatisfactory and, in many cases, is being outpaced by clinical investigation. There are many additional costimulatory molecules of potential therapeutic value, including newly identified B7 ligand family members (130, 131) as well as, undoubtedly, additional as yet uncharacterized regulatory molecules (132). A summary of the current state of our understanding of the biology of these molecules is described in Table 2.

Deeper understanding of the basic biological roles of costimulatory molecules is critical for the rational development of new immune checkpoint blockade therapies. For example, even as therapies targeting other costimulatory molecules move forward in clinical trials, it remains unclear in several instances what the identity of the associated ligand(s) or receptor(s) is, or even whether the target is the receptor or ligand. More ubiquitously, in most cases the precise molecular mechanisms remain unresolved. In addition to cases in which the biology simply remains unknown, there has also been confusion caused by apparently discordant data within the field. Whether these findings reflect additional as yet unappreciated biological complexity or, alternatively, technical differences in experimental systems remains to be fully resolved.

For example, although major histocompatibility complex II (MHC-II) has been previously reported to be the ligand of the coinhibitory receptor LAG3 (133), LSEctin has also been reported to be an additional ligand (134). LSEctin is expressed by liver and tumor cells and may account for the biological role of LAG3 in CD8 and natural killer (NK) cells, as neither cell type interacts with MHC-II. Even more complexity has

been observed in the context of the coinhibitory receptor TIM3, as four ligands have been reported to date: Galectin-9 (135), PtdSer (136), HMGB1 (137), and CEACAM1 (138). How ligand interactions are regulated, whether they affect each other's binding, and whether each ligand leads to unique downstream signaling events remains unclear. Furthermore, although TIM3 is thought of primarily as a marker of T-cell activation and exhaustion, TIM3 also functions to attenuate NK cell cytotoxicity (139). This observation is conceptually significant beyond its pertinence to TIM3 biology, as it suggests that other costimulatory molecules have biologically significant functions in multiple cell types. VISTA presents yet additional ambiguity, as studies have described it as both a ligand on APCs (with homology to PD-L1) with an unknown receptor (140) and as a receptor on T cells with an unknown ligand (141). Similarly, the biological roles of several B7 ligand family members, including their counterreceptors, remain undetermined. B7-H3 is believed to have both costimulatory and coinhibitory roles, possibly dependent on its expression context, whereas both its receptor and the molecular mechanisms of its posttranscriptional regulation remain unclear (142).

Our understanding of the biological functions of costimulatory molecules has been augmented by preclinical and clinical studies using immunomodulatory agents. For example, TIGIT and PD-1 are coexpressed by human melanoma infiltrating NY-ESO-1-specific CD8 T cells (143), consistent with preclinical findings that dual blockade can enhance tumor-infiltrating CD8 T-cell effector function and tumor rejection (144). These findings are consistent with prior observations that TIGIT is induced upon activation and regulates TCR activation pathways in a cell-intrinsic manner (145). These observations suggest that TIGIT and PD-1 blockade may act in *cis*, but through additive mechanisms enhancing T-cell activity. Mechanistic studies have also revealed potential combinatorial strategies to target other nonredundant aspects (e.g., tissue site of action, immune cell population, and biological process) of the cancer-immunity life cycle (51). Analysis of clinical samples reveals that VISTA is expressed primarily on M2 macrophages following ipilimumab treatment in the context of prostate cancer (146). In addition, VISTA and PD-1/PD-L1 have been shown to have nonredundant inhibitory effects on T cells (147). Engagement of innate immunity represents another aspect that can be leveraged to develop effective antitumor immunity. For example, treatment with CD40 agonistic antibodies enhances APC function and, together with chemotherapy, is able to induce effective T cell-dependent immune responses to immunologically “cold” tumors (148, 149). Such studies that identify potential therapies with nonredundant, and ideally synergistic, mechanisms of action will be critical in guiding rational design of combination therapies.

CONCLUDING REMARKS

Here, we have reviewed the current understanding of the biological functions of T-cell costimulatory molecules and the mechanisms through which blockade of these molecules can induce tumor rejection. We have largely focused on CTLA4 and PD-1 immune checkpoint blockade, as well as additional costimulatory molecules of therapeutic interest. Much remains to be understood in how CTLA4, PD-1, and other costimulatory molecules actually attenuate T-cell

Table 2. Summary of the biological and molecular functions of T-cell costimulatory molecules

Molecule	Ligand(s)	Receptor expression pattern	Biological function	Molecular function	References
Coinhibitory					
CTLA4	B7-1 (CD80), B7-2 (CD86)	Activated T cells, Treg	Negative T-cell costimulation (primarily at priming); prevent tonic signaling and/or attenuate high-affinity clones	Competitive inhibition of CD28 costimulation (binding of B7-1 and B7-2)	(8, 10-12, 38, 157-161)
PD-1	PD-L1, PD-L2	Activated T cells, NK cells, NKT cells, B cells, macrophages, subsets of DC; as a result of inflammation	Negative T-cell costimulation (primarily in periphery); attenuate peripheral activity, preserve T-cell function in the context of chronic antigen	Attenuate proximal TCR signaling, attenuate CD28 signaling	(32-35, 38, 39, 53, 100, 162-165)
PD-L1	PD-1, B7-1 (CD80)	Inducible in DC, monocytes, macrophages, mast cells, T cells, B cells, NK cells	Attenuate T-cell activity in inflamed peripheral tissues	PD-1 ligation; cell-intrinsic mechanism unclear	(33, 34, 102)
LAG3	MHC-II, LSECtin	Activated CD4 and CD8 T cells, NK cells, Treg	Negative regulator of T-cell expansion; control T-cell homeostasis; DC activation	Competitive binding to MHC-II; proximal LSECtin mechanism unknown	(133, 134, 166-170)
TIM3	Galectin-9, PtdSer, HMGB1, CEACAM-1	Th1 CD4 and Tc1 CD8, Treg, DC, NK cells, monocytes	Negative regulation of Type 1 immunity; maintain peripheral tolerance	Negative regulation of proximal TCR components; differences between ligands unclear	(135-139, 171)
TIGIT	PVR (CD155), PVRL2 (CD112)	CD4 and CD8, Treg, TFH, NK cells	Negative regulation of T-cell activity; DC tolerization	Competitive inhibition of DNAM1 (CD226) costimulation (binding of PVR), binding of DNAM1 in cis; cell-intrinsic ITIM-negative signaling	(144, 145, 172-176)
VISTA	Counter-receptor unknown	T cells and activated Treg, myeloid cells, mature APC	Negative regulation of T-cell activity; suppression of CD4 T cells	Increase threshold for TCR signaling, induce FOXP3 synthesis; proximal signaling unknown	(140, 141, 146, 147, 177, 178)
Costimulatory					
ICOS	ICOSL	Activated T cells, B cells, ILC2	Positive costimulation; Type I and II immune responses; Treg maintenance; TFH differentiation	p50 PI3K recruitment (AKT signaling); enhance calcium signaling (PLC γ)	(179-186)
OX40	OX40L	Activated T cells, Treg, NK cells, NKT cells, neutrophils	Sustain and enhance CD4 T-cell responses; role in CD8 T cells and Tregs	Regulation of BCL2/XL (survival); enhance PI3K/AKT signaling	(187-193)
GITR	GITRL	Activated T cells, Treg, B cells, NK cells, macrophages	Inhibition of Tregs; costimulation of activated T cells, NK cell activation	Signal through TRAF5	(194-200)
4-1BB (CD137)	4-1BBL	Activated T cells, Treg, NK cells, monocytes, DC, B cells	Positive T-cell costimulation; DC activation	Signal through TRAF1, TRAF2	(201-205)
CD40	CD40L	APCs, B cells, monocytes, nonhematopoietic cells (e.g., fibroblasts, endothelial cells)	APC licensing	Signal through TRAF2, 3, 5, 6; TRAF-independent mechanisms?	(206-209)
CD27	CD70	CD4 and CD8 T cells, B cells, NK cells	Lymphocyte and NK cell costimulation; generation of T-cell memory	Signal through TRAF2, TRAF5	(210-214)

NOTE: A summary of the ligands, immunologic expression pattern, biological function, and molecular mechanisms is presented for selected costimulatory and coinhibitory receptors. Molecular functions (i.e., downstream signaling) reflect predominant currently known mechanisms, but additional mechanisms are likely to contribute significantly.

Abbreviations: NK, natural killer; NKT, natural killer T cell; TFH, T follicular helper; TRAF, tumor necrosis factor receptor-associated factors; DC, dendritic cell.

activation at the molecular, cellular, and physiologic levels. Such mechanistic insight into the biological functions of these molecules will be critical for the development of new approaches and continued improvement of immunotherapeutic strategies. Moving forward, it is likely that combinatorial therapies, utilizing one or more immunotherapies, will become standard of care for a wide breadth of tumor types. Fundamental investigation and understanding of the underlying biology are likely to reveal additional potent biological variables that we have yet to appreciate.

A critical open question is the degree to which the manifestation of irAEs is functionally and mechanistically associated with therapeutic efficacy. If distinct mechanisms underlie these biological responses, a tantalizing possibility is that mechanisms underlying efficacy and irAEs may be able to be engaged separately. Understanding the etiology of irAEs will be even more important in the context of combination therapies, which, at least in the context of anti-CTLA4 and anti-PD-1, have higher rates of irAEs than monotherapies. The safety profile of combination ipilimumab plus nivolumab therapy has been previously reviewed (150). Although most irAEs associated with checkpoint blockade therapy do not reflect the induction of autoimmunity, emerging evidence indicates that autoimmune conditions such as type 1 diabetes and myocarditis can develop at very low frequencies. Fulminant myocarditis has been reported as a potential rare adverse event due to anti-PD-1 monotherapy and combination CTLA4 and PD-1 blockade (151, 152). The development of such rare autoimmune adverse events will become even more relevant as checkpoint blockade therapies are utilized in neoadjuvant (prior to surgery) and adjuvant (following surgery) clinical settings. Understanding how specific immune checkpoint blockade therapies modulate the T-cell repertoire and T-cell function will be essential for distinguishing mechanisms that underlie therapeutic efficacy and irAEs. Indeed, distinct immunologic profiles are associated with colitis induced by anti-CTLA4 and anti-PD-1 therapy (153). One of the key limitations that is currently hampering efforts to understand the manifestation of irAEs is the lack of appropriate preclinical animal models. The development of animal models that faithfully recapitulate irAEs is greatly needed to enable mechanistic investigation of immune checkpoint blockade-associated irAEs.

Of central importance to the mechanisms of action of immune checkpoint blockade therapy is understanding what properties define the antigens that are actually being recognized and mediating tumor rejection. It has been observed that the T-cell repertoire broadens following anti-CTLA4 therapy in patients with melanoma (78). Conversely, response to PD-1 may correlate with reduced intratumoral T-cell clonality (89). This apparent contradiction may reflect observations that tumor regression is often mediated by a small number of dominant neoepitopes (99, 154). Indeed, conservation of abundant clonotypes is associated with better clinical response to anti-CTLA4 therapy (79). Relatedly, PD-1 blockade agents (nivolumab and pembrolizumab) have remarkable efficacy in mismatch repair-deficient and microsatellite instability-high adult and pediatric tumors (83, 84, 155). In addition to these therapies receiving the first tumor tissue-agnostic FDA approval, this is significant because it provides

an example of how mechanistic understanding can identify patient populations likely to benefit from immunotherapeutic approaches. It is important, however, to point out that neoantigen burden represents only one mechanism through which tumors can be recognized by the immune system. Based on the correlation between response rates to anti-PD-1 therapies and TMB across tumor types, it has been estimated that 55% of the variation in therapeutic efficacy can be explained by TMB (87). It is of critical importance to understand additional biological properties, tumor-intrinsic or host-derived, that are significant modulators of therapeutic response. For example, recent studies reveal that genomic lesions in a chromatin remodeling complex component are associated with response to checkpoint blockade (156), providing potential alternative mechanisms of immune recognition.

Immunotherapy has ushered cancer treatment into a new era. In order to translate the progress and success to additional tumor types and to increase the proportion of patients that attain durable responses, we must continue to strive to understand the underlying biological mechanisms. In this review we have highlighted known mechanisms of anti-CTLA4 and anti-PD-1 immune checkpoint inhibitors, but these are by no means complete. There are surely additional surprises awaiting us as we move forward, expanding our understanding of the immune system and its role throughout tumor progression.

Disclosure of Potential Conflicts of Interest

J.P. Allison has ownership interest (including stock, patents, etc.) in Jounce Therapeutics, Forty Seven, ImaginAb, Marker Therapeutics, Tvardi, Constellation, Neon Therapeutics, Apricity, BioAtla, and Polaris, and is a consultant/advisory board member for Jounce Therapeutics, Forty Seven, ImaginAb, Marker Therapeutics, Tvardi, Amgen, Oncolytics, Pieris, Neon Therapeutics, Apricity, BioAtla, and Polaris. No potential conflicts of interest were disclosed by the other authors.

Acknowledgments

We apologize to the many authors and colleagues whose works are not cited due to limited space. We thank J. Mancuso for critical reading of the manuscript and members of the Allison lab for helpful discussions. We also thank Sergio Quezada for helpful discussions. This work was supported by a grant from Cancer Prevention and Research in Texas to J.P. Allison (R1203). S.C. Wei is an MD Anderson Cancer Center Odyssey postdoctoral fellow. Funding as an Odyssey Fellow was supported by the Odyssey Program at The University of Texas MD Anderson Cancer Center. J.P. Allison is a Co-Director of the Parker Institute for Cancer Immunotherapy. J.P. Allison is a cofounder of Jounce and Neon Therapeutics and has received royalties from intellectual property licensed to BMS and Merck.

Received April 4, 2018; revised June 4, 2018; accepted July 11, 2018; published first August 16, 2018.

REFERENCES

1. Sharma P, Allison JP. The future of immune checkpoint therapy. *Science* 2015;348:56–61.
2. Topalian SL, Drake CG, Pardoll DM. Immune checkpoint blockade: a common denominator approach to cancer therapy. *Cancer Cell* 2015;27:450–61.
3. Pardoll DM. The blockade of immune checkpoints in cancer immunotherapy. *Nat Rev Cancer* 2012;12:252–64.
4. Dunn GP, Old LJ, Schreiber RD. The three Es of cancer immunotherapy. *Ann Rev Immunol* 2004;22:329–60.

5. Tang J, Shalabi A, Hubbard-Lucey VM. Comprehensive analysis of the clinical immuno-oncology landscape. *Ann Oncol* 2018;29:84-91.
6. Sharma P, Hu-Lieskovan S, Wargo JA, Ribas A. Primary, adaptive, and acquired resistance to cancer immunotherapy. *Cell* 2017;168:707-23.
7. Ribas A, Wolchok JD. Cancer immunotherapy using checkpoint blockade. *Science* 2018;359:1350-5.
8. Walunas TL, Lenschow DJ, Bakker CY, Linsley PS, Freeman GJ, Green JM, et al. CTLA-4 can function as a negative regulator of T cell activation. *Immunity* 1994;1:405-13.
9. Brunner MC, Chambers CA, Chan FK, Hanke J, Winoto A, Allison JP. CTLA-4-Mediated inhibition of early events of T cell proliferation. *J Immunol* 1999;162:5813-20.
10. Linsley PS, Greene JL, Brady W, Bajorath J, Ledbetter JA, Peach R. Human B7-1 (CD80) and B7-2 (CD86) bind with similar avidities but distinct kinetics to CD28 and CTLA-4 receptors. *Immunity* 1994;1:793-801.
11. Linsley PS, Brady W, Urnes M, Grosmaire LS, Damle NK, Ledbetter JA. CTLA-4 is a second receptor for the B cell activation antigen B7. *J Exp Med* 1991;174:561-9.
12. van der Merwe PA, Bodian DL, Daenke S, Linsley P, Davis SJ. CD80 (B7-1) binds both CD28 and CTLA-4 with a low affinity and very fast kinetics. *J Exp Med* 1997;185:393-403.
13. Lanier LL, O'Fallon S, Somoza C, Phillips JH, Linsley PS, Okumura K, et al. CD80 (B7) and CD86 (B70) provide similar costimulatory signals for T cell proliferation, cytokine production, and generation of CTL. *J Immunol* 1995;154:97-105.
14. Egen JG, Allison JP. Cytotoxic T lymphocyte antigen-4 accumulation in the immunological synapse is regulated by TCR signal strength. *Immunity* 2002;16:23-35.
15. Pentcheva-Hoang T, Egen JG, Wojnoonski K, Allison JP. B7-1 and B7-2 selectively recruit CTLA-4 and CD28 to the immunological synapse. *Immunity* 2004;21:401-13.
16. Kane LP, Andres PG, Howland KC, Abbas AK, Weiss A. Akt provides the CD28 costimulatory signal for up-regulation of IL-2 and IFN-gamma but not TH2 cytokines. *Nat Immunol* 2001;2:37-44.
17. Pages F, Ragueneau M, Rottapel R, Truneh A, Nunes J, Imbert J, et al. Binding of phosphatidylinositol-3-OH kinase to CD28 is required for T-cell signalling. *Nature* 1994;369:327-9.
18. Chambers CA, Cado D, Truong T, Allison JP. Thymocyte development is normal in CTLA-4-deficient mice. *Proc Natl Acad Sci U S A* 1997;94:296-301.
19. Waterhouse P, Penninger JM, Timms E, Wakeham A, Shahinian A, Lee KP, et al. Lymphoproliferative disorders with early lethality in mice deficient in Ctl-4. *Science* 1995;270:985-8.
20. Tivol EA, Borriello F, Schweitzer AN, Lynch WP, Bluestone JA, Sharpe AH. Loss of CTLA-4 leads to massive lymphoproliferation and fatal multiorgan tissue destruction, revealing a critical negative regulatory role of CTLA-4. *Immunity* 1995;3:541-7.
21. Bachmann MF, Kohler G, Ecabert B, Mak TW, Kopf M. Cutting edge: lymphoproliferative disease in the absence of CTLA-4 is not T cell autonomous. *J Immunol* 1999;163:1128-31.
22. Friedline RH, Brown DS, Nguyen H, Kornfeld H, Lee J, Zhang Y, et al. CD4+ regulatory T cells require CTLA-4 for the maintenance of systemic tolerance. *J Exp Med* 2009;206:421-34.
23. Read S, Greenwald R, Izcue A, Robinson N, Mandelbrot D, Francisco L, et al. Blockade of CTLA-4 on CD4+CD25+ regulatory T cells abrogates their function in vivo. *J Immunol* 2006;177:4376-83.
24. Wing K, Onishi Y, Prieto-Martin P, Yamaguchi T, Miyara M, Fehervari Z, et al. CTLA-4 control over Foxp3+ regulatory T cell function. *Science* 2008;322:271-5.
25. Jain N, Nguyen H, Chambers C, Kang J. Dual function of CTLA-4 in regulatory T cells and conventional T cells to prevent multiorgan autoimmunity. *Proc Natl Acad Sci U S A* 2010;107:1524-8.
26. Corse E, Allison JP. Cutting edge: CTLA-4 on effector T cells inhibits in trans. *J Immunol* 2012;189:1123-7.
27. Qureshi OS, Zheng Y, Nakamura K, Attridge K, Manzotti C, Schmidt EM, et al. Trans-endocytosis of CD80 and CD86: a molecular basis for the cell-extrinsic function of CTLA-4. *Science* 2011;332:600-3.
28. Paterson AM, Lovitch SB, Sage PT, Juneja VR, Lee Y, Trombly JD, et al. Deletion of CTLA-4 on regulatory T cells during adulthood leads to resistance to autoimmunity. *J Exp Med* 2015;212:1603-21.
29. Doyle AM, Mullen AC, Villarino AV, Hutchins AS, High FA, Lee HW, et al. Induction of cytotoxic T lymphocyte antigen 4 (CTLA-4) restricts clonal expansion of helper T cells. *J Exp Med* 2001;194:893-902.
30. Egen JG, Kuhns MS, Allison JP. CTLA-4: new insights into its biological function and use in tumor immunotherapy. *Nat Immunol* 2002;3:611-8.
31. Lenschow DJ, Walunas TL, Bluestone JA. CD28/B7 system of T cell costimulation. *Ann Rev Immunol* 1996;14:233-58.
32. Latchman Y, Wood CR, Chernova T, Chaudhary D, Borde M, Chernova I, et al. PD-L2 is a second ligand for PD-1 and inhibits T cell activation. *Nat Immunol* 2001;2:261-8.
33. Freeman GJ, Long AJ, Iwai Y, Bourque K, Chernova T, Nishimura H, et al. Engagement of the PD-1 immunoinhibitory receptor by a novel B7 family member leads to negative regulation of lymphocyte activation. *J Exp Med* 2000;192:1027-34.
34. Dong H, Zhu G, Tamada K, Chen L. B7-H1, a third member of the B7 family, co-stimulates T-cell proliferation and interleukin-10 secretion. *Nat Med* 1999;5:1365-9.
35. Agata Y, Kawasaki A, Nishimura H, Ishida Y, Tsubata T, Yagita H, et al. Expression of the PD-1 antigen on the surface of stimulated mouse T and B lymphocytes. *Int Immunol* 1996;8:765-72.
36. Keir ME, Liang SC, Guleria I, Latchman YE, Qipo A, Albacker LA, et al. Tissue expression of PD-L1 mediates peripheral T cell tolerance. *J Exp Med* 2006;203:883-95.
37. Yokosuka T, Takamatsu M, Kobayashi-Imanishi W, Hashimoto-Tane A, Azuma M, Saito T. Programmed cell death 1 forms negative costimulatory microclusters that directly inhibit T cell receptor signaling by recruiting phosphatase SHP2. *J Exp Med* 2012;209:1201-17.
38. Parry RV, Chemnitz JM, Frauwirth KA, Lanfranco AR, Braunstein I, Kobayashi SV, et al. CTLA-4 and PD-1 receptors inhibit T-cell activation by distinct mechanisms. *Mol Cell Biol* 2005;25:9543-53.
39. Hui E, Cheung J, Zhu J, Su X, Taylor MJ, Wallweber HA, et al. T cell costimulatory receptor CD28 is a primary target for PD-1-mediated inhibition. *Science* 2017;355:1428-33.
40. Rota G, Niogret C, Dang AT, Barros CR, Fonta NP, Alfei F, et al. Shp-2 is dispensable for establishing T cell exhaustion and for PD-1 signaling in vivo. *Cell reports* 2018;23:39-49.
41. Nishimura H, Nose M, Hiai H, Minato N, Honjo T. Development of lupus-like autoimmune diseases by disruption of the PD-1 gene encoding an ITIM motif-carrying immunoreceptor. *Immunity* 1999;11:141-51.
42. Nishimura H, Okazaki T, Tanaka Y, Nakatani K, Hara M, Matsumori A, et al. Autoimmune dilated cardiomyopathy in PD-1 receptor-deficient mice. *Science* 2001;291:319-22.
43. Bengsch B, Johnson AL, Kurachi M, Odorizzi PM, Pauken KE, Attanasio J, et al. Bioenergetic insufficiencies due to metabolic alterations regulated by the inhibitory receptor PD-1 are an early driver of CD8(+) T cell exhaustion. *Immunity* 2016;45:358-73.
44. Patsoukis N, Bardhan K, Chatterjee P, Sari D, Liu B, Bell LN, et al. PD-1 alters T-cell metabolic reprogramming by inhibiting glycolysis and promoting lipolysis and fatty acid oxidation. *Nat Commun* 2015;6:6692.
45. Buck MD, O'Sullivan D, Klein Geltink RI, Curtis JD, Chang CH, Sanin DE, et al. Mitochondrial dynamics controls T cell fate through metabolic programming. *Cell* 2016;166:63-76.
46. Wherry EJ, Ha SJ, Kaech SM, Haining WN, Sarkar S, Kalia V, et al. Molecular signature of CD8+ T cell exhaustion during chronic viral infection. *Immunity* 2007;27:670-84.
47. Pauken KE, Sammons MA, Odorizzi PM, Manne S, Godec J, Khan O, et al. Epigenetic stability of exhausted T cells limits durability of reinvigoration by PD-1 blockade. *Science* 2016;354:1160-5.
48. Philip M, Fairchild L, Sun L, Horste EL, Camara S, Shakiba M, et al. Chromatin states define tumour-specific T cell dysfunction and reprogramming. *Nature* 2017;545:452-6.
49. Kortlever RM, Sodikin NM, Wilson CH, Burkhardt DL, Pellegrinet L, Brown Swigart L, et al. Myc cooperates with ras by programming inflammation and immune suppression. *Cell* 2017;171:1301-15 e1314.

50. Kleffel S, Posch C, Barthel SR, Mueller H, Schlapbach C, Guenova E, et al. Melanoma cell-intrinsic PD-1 receptor functions promote tumor growth. *Cell* 2015;162:1242–56.
51. Chen DS, Mellman I. Oncology meets immunology: the cancer-immunity cycle. *Immunity* 2013;39:1–10.
52. Wherry EJ. T cell exhaustion. *Nat Immunol* 2011;12:492–9.
53. Barber DL, Wherry EJ, Masopust D, Zhu B, Allison JP, Sharpe AH, et al. Restoring function in exhausted CD8 T cells during chronic viral infection. *Nature* 2006;439:682–7.
54. Kamphorst AO, Wieland A, Nasti T, Yang S, Zhang R, Barber DL, et al. Rescue of exhausted CD8 T cells by PD-1-targeted therapies is CD28-dependent. *Science* 2017;355:1423–7.
55. Ramagopal UA, Liu W, Garrett-Thomson SC, Bonanno JB, Yan Q, Srinivasan M, et al. Structural basis for cancer immunotherapy by the first-in-class checkpoint inhibitor ipilimumab. *Proc Natl Acad Sci U S A* 2017;114:E4223–32.
56. Dieu-Nosjean MC, Antoine M, Danel C, Heudes D, Wislez M, Poulot V, et al. Long-term survival for patients with non-small-cell lung cancer with intratumoral lymphoid structures. *J Clin Oncol* 2008;26:4410–7.
57. Engelhard VH, Rodriguez AB, Mauldin IS, Woods AN, Peske JD, Slingluff CL Jr. Immune cell infiltration and tertiary lymphoid structures as determinants of antitumor immunity. *J Immunol* 2018;200:432–42.
58. Joshi NS, Akama-Garren EH, Lu Y, Lee DY, Chang GP, Li A, et al. Regulatory T cells in tumor-associated tertiary lymphoid structures suppress anti-tumor T cell responses. *Immunity* 2015;43:579–90.
59. Fehlings M, Simoni Y, Penny HL, Becht E, Loh CY, Gubin MM, et al. Checkpoint blockade immunotherapy reshapes the high-dimensional phenotypic heterogeneity of murine intratumoral neoantigen-specific CD8(+) T cells. *Nat Commun* 2017;8:562.
60. Wei SC, Levine JH, Cogdill AP, Zhao Y, Anang NAS, Andrews MC, et al. Distinct cellular mechanisms underlie anti-CTLA-4 and anti-PD-1 checkpoint blockade. *Cell* 2017;170:1120–33 e1117.
61. Liakou CI, Kamat A, Tang DN, Chen H, Sun J, Troncoso P, et al. CTLA-4 blockade increases IFN γ -producing CD4+ICOShi cells to shift the ratio of effector to regulatory T cells in cancer patients. *Proc Natl Acad Sci U S A* 2008;105:14987–92.
62. Carthon BC, Wolchok JD, Yuan J, Kamat A, Ng Tang DS, Sun J, et al. Preoperative CTLA-4 blockade: tolerability and immune monitoring in the setting of a presurgical clinical trial. *Clin Cancer Res* 2010;16:2861–71.
63. Chen H, Liakou CI, Kamat A, Pettaway C, Ward JF, Tang DN, et al. Anti-CTLA-4 therapy results in higher CD4+ICOShi T cell frequency and IFN- γ levels in both nonmalignant and malignant prostate tissues. *Proc Natl Acad Sci U S A* 2009;106:2729–34.
64. Chaput N, Lepage P, Coutzac C, Soularue E, Le Roux K, Monot C, et al. Baseline gut microbiota predicts clinical response and colitis in metastatic melanoma patients treated with ipilimumab. *Ann Oncol* 2017;28:1368–79.
65. Vonderheide RH, LoRusso PM, Khalil M, Gartner EM, Khaira D, Soulieres D, et al. Tremelimumab in combination with exemestane in patients with advanced breast cancer and treatment-associated modulation of inducible costimulator expression on patient T cells. *Clin Cancer Res* 2010;16:3485–94.
66. Ng Tang D, Shen Y, Sun J, Wen S, Wolchok JD, Yuan J, et al. Increased frequency of ICOS+ CD4 T cells as a pharmacodynamic biomarker for anti-CTLA-4 therapy. *Cancer Immunol Res* 2013;1:229–34.
67. Simpson TR, Li F, Montalvo-Ortiz W, Sepulveda MA, Bergerhoff K, Arce F, et al. Fc-dependent depletion of tumor-infiltrating regulatory T cells co-defines the efficacy of anti-CTLA-4 therapy against melanoma. *J Exp Med* 2013;210:1695–710.
68. Selby MJ, Engelhardt JJ, Quigley M, Henning KA, Chen T, Srinivasan M, et al. Anti-CTLA-4 antibodies of IgG2a isotype enhance antitumor activity through reduction of intratumoral regulatory T cells. *Cancer Immunol Res* 2013;1:32–42.
69. Bulliard Y, Jolicoeur R, Windman M, Rue SM, Ettenberg S, Knee DA, et al. Activating Fc gamma receptors contribute to the antitumor activities of immunoregulatory receptor-targeting antibodies. *J Exp Med* 2013;210:1685–93.
70. Du X, Tang F, Liu M, Su J, Zhang Y, Wu W, et al. A reappraisal of CTLA-4 checkpoint blockade in cancer immunotherapy. *Cell Res* 2018;28:416–32.
71. Peggs KS, Quezada SA, Chambers CA, Korman AJ, Allison JP. Blockade of CTLA-4 on both effector and regulatory T cell compartments contributes to the antitumor activity of anti-CTLA-4 antibodies. *J Exp Med* 2009;206:1717–25.
72. Romano E, Kusio-Kobialka M, Foukas PG, Baumgaertner P, Meyer C, Ballabeni P, et al. Ipilimumab-dependent cell-mediated cytotoxicity of regulatory T cells ex vivo by nonclassical monocytes in melanoma patients. *Proc Natl Acad Sci U S A* 2015;112:6140–5.
73. Arce Vargas F, Furness AJS, Litchfield K, Joshi K, Rosenthal R, Ghorani E, et al. Fc effector function contributes to the activity of human anti-CTLA-4 antibodies. *Cancer Cell* 2018;33:649–63.e4.
74. Bruhns P, Iannascoli B, England P, Mancardi DA, Fernandez N, Jorieux S, et al. Specificity and affinity of human Fc γ receptors and their polymorphic variants for human IgG subclasses. *Blood* 2009;113:3716–25.
75. Ribas A, Kefford R, Marshall MA, Punt CJ, Haanen JB, Marmol M, et al. Phase III randomized clinical trial comparing tremelimumab with standard-of-care chemotherapy in patients with advanced melanoma. *J Clin Oncol* 2013;31:616–22.
76. Eroglu Z, Kim DW, Wang X, Camacho LH, Chmielowski B, Seja E, et al. Long term survival with cytotoxic T lymphocyte-associated antigen 4 blockade using tremelimumab. *Eur J Cancer* 2015;51:2689–97.
77. Schadendorf D, Hodi FS, Robert C, Weber JS, Margolin K, Hamid O, et al. Pooled analysis of long-term survival data from phase II and phase III trials of ipilimumab in unresectable or metastatic melanoma. *J Clin Oncol* 2015;33:1889–94.
78. Robert L, Tsoi J, Wang X, Emerson R, Homet B, Chodon T, et al. CTLA4 blockade broadens the peripheral T-cell receptor repertoire. *Clin Cancer Res* 2014;20:2424–32.
79. Cha E, Klinger M, Hou Y, Cummings C, Ribas A, Faham M, et al. Improved survival with T cell clonotype stability after anti-CTLA-4 treatment in cancer patients. *Sci Translat Med* 2014;6:238ra270.
80. Kvistborg P, Philips D, Kelderman S, Hageman L, Ottensmeier C, Joseph-Pietras D, et al. Anti-CTLA-4 therapy broadens the melanoma-reactive CD8+ T cell response. *Sci Translat Med* 2014;6:254ra128.
81. Oh DY, Cham J, Zhang L, Fong G, Kwek SS, Klinger M, et al. Immune toxicities elicited by CTLA-4 blockade in cancer patients are associated with early diversification of the T-cell repertoire. *Cancer Res* 2017;77:1322–30.
82. Hellmann MD, Nathanson T, Rizvi H, Creelan BC, Sanchez-Vega F, Ahuja A, et al. Genomic features of response to combination immunotherapy in patients with advanced non-small-cell lung cancer. *Cancer Cell* 2018;33:843–52.e4.
83. Le DT, Uram JN, Wang H, Bartlett BR, Kemberling H, Eyring AD, et al. PD-1 blockade in tumors with mismatch-repair deficiency. *N Engl J Med* 2015;372:2509–20.
84. Le DT, Durham JN, Smith KN, Wang H, Bartlett BR, Aulakh LK, et al. Mismatch repair deficiency predicts response of solid tumors to PD-1 blockade. *Science* 2017;357:409–13.
85. Snyder A, Makarov V, Merghoub T, Yuan J, Zaretsky JM, Desrichard A, et al. Genetic basis for clinical response to CTLA-4 blockade in melanoma. *N Engl J Med* 2014;371:2189–99.
86. Van Allen EM, Miao D, Schilling B, Shukla SA, Blank C, Zimmer L, et al. Genomic correlates of response to CTLA-4 blockade in metastatic melanoma. *Science* 2015;350:207–11.
87. Yarchoan M, Hopkins A, Jaffee EM. Tumor mutational burden and response rate to PD-1 inhibition. *N Engl J Med* 2017;377:2500–1.
88. Herbst RS, Soria JC, Kowanetz M, Fine GD, Hamid O, Gordon MS, et al. Predictive correlates of response to the anti-PD-L1 antibody MPDL3280A in cancer patients. *Nature* 2014;515:563–7.
89. Tumei PC, Harview CL, Yearley JH, Shintaku IP, Taylor EJ, Robert L, et al. PD-1 blockade induces responses by inhibiting adaptive immune resistance. *Nature* 2014;515:568–71.

90. Im SJ, Hashimoto M, Gerner MY, Lee J, Kissick HT, Burger MC, et al. Defining CD8+ T cells that provide the proliferative burst after PD-1 therapy. *Nature* 2016;537:417–21.
91. Huang AC, Postow MA, Orlowski RJ, Mick R, Bengsch B, Manne S, et al. T-cell invigoration to tumour burden ratio associated with anti-PD-1 response. *Nature* 2017;545:60–5.
92. Forde PM, Chaft JE, Smith KN, Anagnostou V, Cottrell TR, Hellmann MD, et al. Neoadjuvant PD-1 blockade in resectable lung cancer. *N Engl J Med* 2018;378:1976–86.
93. Ghoneim HE, Fan Y, Moustaki A, Abdelsamed HA, Dash P, Dogra P, et al. De novo epigenetic programs inhibit PD-1 blockade-mediated T cell rejuvenation. *Cell* 2017;170:142–57 e119.
94. Sen DR, Kaminski J, Barnitz RA, Kurachi M, Gerdemann U, Yates KB, et al. The epigenetic landscape of T cell exhaustion. *Science* 2016;354:1165–9.
95. Bengsch B, Ohtani T, Khan O, Setty M, Manne S, O'Brien S, et al. Epigenomic-guided mass cytometry profiling reveals disease-specific features of exhausted CD8 T cells. *Immunity* 2018;48:1029–45 e1025.
96. Spitzer MH, Carmi Y, Reticker-Flynn NE, Kwek SS, Madhiredy D, Martins MM, et al. Systemic immunity is required for effective cancer immunotherapy. *Cell* 2017;168:487–502 e415.
97. Iijima N, Iwasaki A. Access of protective antiviral antibody to neuronal tissues requires CD4 T-cell help. *Nature* 2016;533:552–6.
98. Nakanishi Y, Lu B, Gerard C, Iwasaki A. CD8(+) T lymphocyte mobilization to virus-infected tissue requires CD4(+) T-cell help. *Nature* 2009;462:510–3.
99. Gubin MM, Zhang X, Schuster H, Caron E, Ward JP, Noguchi T, et al. Checkpoint blockade cancer immunotherapy targets tumour-specific mutant antigens. *Nature* 2014;515:577–81.
100. Loke P, Allison JP. PD-L1 and PD-L2 are differentially regulated by Th1 and Th2 cells. *Proc Natl Acad Sci U S A* 2003;100:5336–41.
101. Dahan R, Segal E, Engelhardt J, Selby M, Korman AJ, Ravetch JV. FcγR3 modulate the anti-tumor activity of antibodies targeting the PD-1/PD-L1 Axis. *Cancer Cell* 2015;28:285–95.
102. Butte MJ, Keir ME, Phamduy TB, Sharpe AH, Freeman GJ. Programmed death-1 ligand 1 interacts specifically with the B7-1 costimulatory molecule to inhibit T cell responses. *Immunity* 2007;27:111–22.
103. Tang H, Liang Y, Anders RA, Taube JM, Qiu X, Mulgaonkar A, et al. PD-L1 on host cells is essential for PD-L1 blockade-mediated tumor regression. *J Clin Invest* 2018;128:580–88.
104. Lin H, Wei S, Hurt EM, Green MD, Zhao L, Vatan L, et al. Host expression of PD-L1 determines efficacy of PD-L1 pathway blockade-mediated tumor regression. *J Clin Invest* 2018;128(2):805–15.
105. Juneja VR, McGuire KA, Manguso RT, LaFleur MW, Collins N, Haining WN, et al. PD-L1 on tumor cells is sufficient for immune evasion in immunogenic tumors and inhibits CD8 T cell cytotoxicity. *J Exp Med* 2017;214:895–904.
106. Postow MA, Chesney J, Pavlick AC, Robert C, Grossmann K, McDermott D, et al. Nivolumab and ipilimumab versus ipilimumab in untreated melanoma. *N Engl J Med* 2015;372:2006–17.
107. Larkin J, Chiarion-Sileni V, Gonzalez R, Grob JJ, Cowey CL, Lao CD, et al. Combined nivolumab and ipilimumab or monotherapy in untreated melanoma. *N Engl J Med* 2015;373:23–34.
108. Wolchok JD, Kluger H, Callahan MK, Postow MA, Rizvi NA, Lesokhin AM, et al. Nivolumab plus ipilimumab in advanced melanoma. *N Engl J Med* 2013;369:122–33.
109. Curran MA, Montalvo W, Yagita H, Allison JP. PD-1 and CTLA-4 combination blockade expands infiltrating T cells and reduces regulatory T and myeloid cells within B16 melanoma tumors. *Proc Natl Acad Sci U S A* 2010;107:4275–80.
110. Postow MA, Larkin J, Wolchok J, Chiarion-Sileni V, Hodi FS, Rutkowski P, et al. 32nd annual meeting and pre-conference programs of the society for immunotherapy of cancer (SITC 2017): part one. Pooled 3-year overall survival data from phase II and phase III trials of nivolumab (NIVO) combined with ipilimumab (IPI) in advanced melanoma. *J Immunother Cancer* 2017;5:86.
111. Motzer RJ, Tannir NM, McDermott DF, Arén Frontera O, Melichar B, Choueiri TK, et al. Nivolumab plus ipilimumab versus sunitinib in advanced renal-cell carcinoma. *N Engl J Med* 2018;378:1277–90.
112. Das R, Verma R, Sznol M, Boddupalli CS, Gettinger SN, Kluger H, et al. Combination therapy with anti-CTLA-4 and anti-PD-1 leads to distinct immunologic changes in vivo. *J Immunol* 2015;194:950–9.
113. Das R, Bar N, Ferreira M, Newman AM, Zhang L, Bailur JK, et al. Early B cell changes predict autoimmunity following combination immune checkpoint blockade. *J Clin Invest* 2018;128:715–20.
114. Kamphorst AO, Pillai RN, Yang S, Nasti TH, Akondy RS, Wieland A, et al. Proliferation of PD-1+ CD8 T cells in peripheral blood after PD-1-targeted therapy in lung cancer patients. *Proc Natl Acad Sci U S A* 2017;114:4993–8.
115. Daud AI, Loo K, Pauli ML, Sanchez-Rodriguez R, Sandoval PM, Taravati K, et al. Tumor immune profiling predicts response to anti-PD-1 therapy in human melanoma. *J Clin Invest* 2016;126:3447–52.
116. Weber JS, Gibney G, Sullivan RJ, Sosman JA, Slingluff CL Jr, Lawrence DP, et al. Sequential administration of nivolumab and ipilimumab with a planned switch in patients with advanced melanoma (CheckMate 064): an open-label, randomised, phase 2 trial. *Lancet Oncol* 2016;17:943–55.
117. Sanchez-Paulete AR, Cueto FJ, Martínez-López M, Labiano S, Morales-Kastresana A, Rodríguez-Ruiz ME, et al. Cancer immunotherapy with immunomodulatory anti-CD137 and anti-PD-1 monoclonal antibodies requires BATF3-Dependent dendritic cells. *Cancer Discov* 2016;6:71–9.
118. Salmon H, Idoyaga J, Rahman A, Leboeuf M, Remark R, Jordan S, et al. Expansion and activation of CD103(+) dendritic cell progenitors at the tumor site enhances tumor responses to therapeutic PD-L1 and BRAF inhibition. *Immunity* 2016;44:924–38.
119. Spranger S, Dai D, Horton B, Gajewski TF. Tumor-residing Batf3 dendritic cells are required for effector T cell trafficking and adoptive T cell therapy. *Cancer Cell* 2017;31:711–23 e714.
120. Roberts EW, Broz ML, Binnewies M, Headley MB, Nelson AE, Wolf DM, et al. Critical role for CD103(+)/CD141(+) dendritic cells bearing CCR7 for tumor antigen trafficking and priming of T cell immunity in melanoma. *Cancer Cell* 2016;30:324–36.
121. Twyman-Saint Victor C, Rech AJ, Maity A, Rengan R, Pauken KE, Stelekati E, et al. Radiation and dual checkpoint blockade activate non-redundant immune mechanisms in cancer. *Nature* 2015;520:373–7.
122. Dovedi SJ, Adlard AL, Lipowska-Bhalla G, McKenna C, Jones S, Cheadle EJ, et al. Acquired resistance to fractionated radiotherapy can be overcome by concurrent PD-L1 blockade. *Cancer Res* 2014;74:5458–68.
123. Demaria S, Kawashima N, Yang AM, Devitt ML, Babb JS, Allison JP, et al. Immune-mediated inhibition of metastases after treatment with local radiation and CTLA-4 blockade in a mouse model of breast cancer. *Clin Cancer Res* 2005;11:728–34.
124. Golden EB, Demaria S, Schiff PB, Chachoua A, Formenti SC. An abscopal response to radiation and ipilimumab in a patient with metastatic non-small cell lung cancer. *Cancer Immunol Res* 2013;1:365–72.
125. De Henau O, Rausch M, Winkler D, Campesato LF, Liu C, Cymerman DH, et al. Overcoming resistance to checkpoint blockade therapy by targeting PI3Kγ in myeloid cells. *Nature* 2016;539:443–7.
126. Sivan A, Corrales L, Hubert N, Williams JB, Aquino-Michaels K, Earley ZM, et al. Commensal Bifidobacterium promotes antitumor immunity and facilitates anti-PD-L1 efficacy. *Science* 2015;350:1084–9.
127. Vetzizou M, Pitt JM, Daillère R, Lepage P, Waldschmitt N, Flament C, et al. Anticancer immunotherapy by CTLA-4 blockade relies on the gut microbiota. *Science* 2015;350:1079–84.
128. Routy B, Le Chatelier E, Derosa L, Duong CPM, Alou MT, Daillère R, et al. Gut microbiome influences efficacy of PD-1-based immunotherapy against epithelial tumors. *Science* 2018;359:91–7.
129. Gopalakrishnan V, Spencer CN, Nezi L, Reuben A, Andrews MC, Karpman TV, et al. Gut microbiome modulates response to anti-PD-1 immunotherapy in melanoma patients. *Science* 2018;359:97–103.
130. Ni L, Dong C. New checkpoints in cancer immunotherapy. *Immunol Rev* 2017;276:52–65.
131. Janakiram M, Shah UA, Liu W, Zhao A, Schoenberg MP, Zang X, et al. The third group of the B7-CD28 immune checkpoint family: HHLA2, TMIGD2, B7x, and B7-H3. *Immunol Rev* 2017;276:26–39.

132. Pentcheva-Hoang T, Corse E, Allison JP. Negative regulators of T-cell activation: potential targets for therapeutic intervention in cancer, autoimmune disease, and persistent infections. *Immunol Rev* 2009;229:67–87.
133. Huard B, Prigent P, Tournier M, Bruniquel D, Triebel F. CD4/major histocompatibility complex class II interaction analyzed with CD4- and lymphocyte activation gene-3 (LAG-3)-Ig fusion proteins. *Eur J Immunol* 1995;25:2718–21.
134. Xu F, Liu J, Liu D, Liu B, Wang M, Hu Z, et al. LSECtin expressed on melanoma cells promotes tumor progression by inhibiting antitumor T-cell responses. *Cancer Res* 2014;74:3418–28.
135. Zhu C, Anderson AC, Schubart A, Xiong H, Imitola J, Khoury SJ, et al. The Tim-3 ligand galectin-9 negatively regulates T helper type 1 immunity. *Nat Immunol* 2005;6:1245–52.
136. DeKruyff RH, Bu X, Ballesteros A, Santiago C, Chim YL, Lee HH, et al. T cell/transmembrane, Ig, and mucin-3 allelic variants differentially recognize phosphatidylserine and mediate phagocytosis of apoptotic cells. *J Immunol* 2010;184:1918–30.
137. Chiba S, Baghdadi M, Akiba H, Yoshiyama H, Kinoshita I, Dosaka-Akita H, et al. Tumor-infiltrating DCs suppress nucleic acid-mediated innate immune responses through interactions between the receptor TIM-3 and the alarmin HMGB1. *Nat Immunol* 2012;13:832–42.
138. Huang YH, Zhu C, Kondo Y, Anderson AC, Gandhi A, Russell A, et al. CEACAM1 regulates TIM-3-mediated tolerance and exhaustion. *Nature* 2015;517:386–90.
139. Ndhlovu LC, Lopez-Vergès S, Barbour JD, Jones RB, Jha AR, Long BR, et al. Tim-3 marks human natural killer cell maturation and suppresses cell-mediated cytotoxicity. *Blood* 2012;119:3734–43.
140. Wang L, Rubinstein R, Lines JL, Wasiuk A, Ahonen C, Guo Y, et al. VISTA, a novel mouse Ig superfamily ligand that negatively regulates T cell responses. *J Exp Med* 2011;208:577–92.
141. Flies DB, Han X, Higuchi T, Zheng L, Sun J, Ye JJ, et al. Coinhibitory receptor PD-1H preferentially suppresses CD4(+) T cell-mediated immunity. *J Clin Invest* 2014;124:1966–75.
142. Schildberg FA, Klein SR, Freeman GJ, Sharpe AH. Coinhibitory Pathways in the B7-CD28 Ligand-Receptor Family. *Immunity* 2016;44:955–72.
143. Chauvin JM, Pagliano O, Fourcade J, Sun Z, Wang H, Sander C, et al. TIGIT and PD-1 impair tumor antigen-specific CD8(+) T cells in melanoma patients. *J Clin Invest* 2015;125:2046–58.
144. Johnston RJ, Comps-Agrar L, Hackney J, Yu X, Huseni M, Yang Y, et al. The immunoreceptor TIGIT regulates antitumor and antiviral CD8(+) T cell effector function. *Cancer Cell* 2014;26:923–37.
145. Joller N, Hafler JP, Brynedal B, Kassam N, Spoerl S, Levin SD, et al. Cutting edge: TIGIT has T cell-intrinsic inhibitory functions. *J Immunol* 2011;186:1338–42.
146. Gao J, Ward JF, Pettaway CA, Shi LZ, Subudhi SK, Vence LM, et al. VISTA is an inhibitory immune checkpoint that is increased after ipilimumab therapy in patients with prostate cancer. *Nat Med* 2017;23:551–5.
147. Liu J, Yuan Y, Chen W, Putra J, Suriawinata AA, Schenk AD, et al. Immune-checkpoint proteins VISTA and PD-1 nonredundantly regulate murine T-cell responses. *Proc Natl Acad Sci U S A* 2015;112:6682–7.
148. Byrne KT, Vonderheide RH. CD40 stimulation obviates innate sensors and drives T cell immunity in cancer. *Cell Rep* 2016;15:2719–32.
149. Vonderheide RH, Glennie MJ. Agonistic CD40 antibodies and cancer therapy. *Clin Cancer Res* 2013;19:1035–43.
150. Sznol M, Ferrucci PF, Hogg D, Atkins MB, Wolter P, Guidoboni M, et al. Pooled analysis safety profile of nivolumab and ipilimumab combination therapy in patients with advanced melanoma. *J Clin Oncol* 2017;35:3815–22.
151. Johnson DB, Balko JM, Compton ML, Chalkias S, Gorham J, Xu Y, et al. Fulminant myocarditis with combination immune checkpoint blockade. *N Engl J Med* 2016;375:1749–55.
152. Moslehi JJ, Salem JE, Sosman JA, Lebrun-Vignes B, Johnson DB. Increased reporting of fatal immune checkpoint inhibitor-associated myocarditis. *Lancet* 2018;391:933.
153. Coutzac C, Adam J, Soularue E, Collins M, Racine A, Mussini C, et al. Colon immune-related adverse events: anti-CTLA-4 and anti-PD-1 blockade induce distinct immunopathological entities. *J Crohns Colitis* 2017;11:1238–46.
154. Matsushita H, Vesely MD, Koboldt DC, Rickert CG, Uppaluri R, Magrini VJ, et al. Cancer exome analysis reveals a T-cell-dependent mechanism of cancer immunoeediting. *Nature* 2012;482:400–4.
155. Overman MJ, McDermott R, Leach JL, Lonardi S, Lenz HJ, Morse MA, et al. Nivolumab in patients with metastatic DNA mismatch repair-deficient or microsatellite instability-high colorectal cancer (CheckMate 142): an open-label, multicentre, phase 2 study. *Lancet Oncol* 2017;18:1182–91.
156. Miao D, Margolis CA, Gao W, Voss MH, Li W, Martini DJ, et al. Genomic correlates of response to immune checkpoint therapies in clear cell renal cell carcinoma. *Science* 2018;359:801–6.
157. Freeman GJ, Gribben JG, Boussiotis VA, Ng JW, Restivo VA Jr, Lombard LA, et al. Cloning of B7-2: a CTLA-4 counter-receptor that costimulates human T cell proliferation. *Science* 1993;262:909–11.
158. Freeman GJ, Borriello F, Hodes RJ, Reiser H, Hathcock KS, Laszlo G, et al. Uncovering of functional alternative CTLA-4 counter-receptor in B7-deficient mice. *Science* 1993;262:907–9.
159. Krummel MF, Allison JP. CD28 and CTLA-4 have opposing effects on the response of T cells to stimulation. *J Exp Med* 1995;182:459–65.
160. Krummel MF, Allison JP. CTLA-4 engagement inhibits IL-2 accumulation and cell cycle progression upon activation of resting T cells. *J Exp Med* 1996;183:2533–40.
161. Lenschow DJ, Su GH, Zuckerman LA, Nabavi N, Jellis CL, Gray GS, et al. Expression and functional significance of an additional ligand for CTLA-4. *Proc Natl Acad Sci U S A* 1993;90:11054–8.
162. Ishida Y, Agata Y, Shibahara K, Honjo T. Induced expression of PD-1, a novel member of the immunoglobulin gene superfamily, upon programmed cell death. *EMBO J* 1992;11:3887–95.
163. Sheppard KA, Fitz LJ, Lee JM, Benander C, George JA, Wooters J, et al. PD-1 inhibits T-cell receptor induced phosphorylation of the ZAP70/CD3zeta signalosome and downstream signaling to PKCtheta. *FEBS Lett* 2004;574:37–41.
164. Tseng SY, Otsuji M, Gorski K, Huang X, Slansky JE, Pai SI, et al. B7-DC, a new dendritic cell molecule with potent costimulatory properties for T cells. *J Exp Med* 2001;193:839–46.
165. Chemnitz JM, Parry RV, Nichols KE, June CH, Riley JL. SHP-1 and SHP-2 associate with immunoreceptor tyrosine-based switch motif of programmed death 1 upon primary human T cell stimulation, but only receptor ligation prevents T cell activation. *J Immunol* 2004;173:945–54.
166. Triebel F, Jitsukawa S, Baixeras E, Roman-Roman S, Genevee C, Viegas-Pequignot E, et al. LAG-3, a novel lymphocyte activation gene closely related to CD4. *J Exp Med* 1990;171:1393–405.
167. Huard B, Mastrangeli R, Prigent P, Bruniquel D, Donini S, El-Tayar N, et al. Characterization of the major histocompatibility complex class II binding site on LAG-3 protein. *Proc Natl Acad Sci U S A* 1997;94:5744–9.
168. Workman CJ, Dugger KJ, Vignali DA. Cutting edge: molecular analysis of the negative regulatory function of lymphocyte activation gene-3. *J Immunol* 2002;169:5392–5.
169. Workman CJ, Vignali DA. Negative regulation of T cell homeostasis by lymphocyte activation gene-3 (CD223). *J Immunol* 2005;174:688–95.
170. Andrae S, Piras F, Burdin N, Triebel F. Maturation and activation of dendritic cells induced by lymphocyte activation gene-3 (CD223). *J Immunol* 2002;168:3874–80.
171. Monney L, Sabatos CA, Gaglia JL, Ryu A, Waldner H, Chernova T, et al. Th1-specific cell surface protein Tim-3 regulates macrophage activation and severity of an autoimmune disease. *Nature* 2002;415:536–41.
172. Stanitsky N, Simic H, Arapovic J, Toporik A, Levy O, Novik A, et al. The interaction of TIGIT with PVR and PVRL2 inhibits human NK cell cytotoxicity. *Proc Natl Acad Sci U S A* 2009;106:17858–63.
173. Yu X, Harden K, Gonzalez LC, Francesco M, Chiang E, Irving B, et al. The surface protein TIGIT suppresses T cell activation by promoting the generation of mature immunoregulatory dendritic cells. *Nat Immunol* 2009;10:48–57.
174. Levin SD, Taft DW, Brandt CS, Bucher C, Howard ED, Chadwick EM, et al. Vstm3 is a member of the CD28 family and an important modulator of T-cell function. *Eur J Immunol* 2011;41:902–15.

175. Boles KS, Vermi W, Facchetti F, Fuchs A, Wilson TJ, Diacovo TG, et al. A novel molecular interaction for the adhesion of follicular CD4 T cells to follicular DC. *Eur J Immunol* 2009;39:695–703.
176. Stanitsky N, Rovis TL, Glasner A, Seidel E, Tsukerman P, Yamin R, et al. Mouse TIGIT inhibits NK-cell cytotoxicity upon interaction with PVR. *Eur J Immunol* 2013;43:2138–50.
177. Lines JL, Pantazi E, Mak J, Sempere LF, Wang L, O'Connell S, et al. VISTA is an immune checkpoint molecule for human T cells. *Cancer Res* 2014;74:1924–32.
178. Wang L, Le Mercier I, Putra J, Chen W, Liu J, Schenk AD, et al. Disruption of the immune-checkpoint VISTA gene imparts a proinflammatory phenotype with predisposition to the development of autoimmunity. *Proc Natl Acad Sci U S A* 2014;111:14846–51.
179. Yoshinaga SK, Whoriskey JS, Khare SD, Sarmiento U, Guo J, Horan T, et al. T-cell co-stimulation through B7RP-1 and ICOS. *Nature* 1999;402:827–32.
180. McAdam AJ, Chang TT, Lumelsky AE, Greenfield EA, Boussiotis VA, Duke-Cohan JS, et al. Mouse inducible costimulatory molecule (ICOS) expression is enhanced by CD28 costimulation and regulates differentiation of CD4+ T cells. *J Immunol* 2000;165:5035–40.
181. Burmeister Y, Lischke T, Dahler AC, Mages HW, Lam KP, Coyle AJ, et al. ICOS controls the pool size of effector-memory and regulatory T cells. *J Immunol* 2008;180:774–82.
182. Fos C, Salles A, Lang V, Carrette F, Audebert S, Pastor S, et al. ICOS ligation recruits the p50alpha PI3K regulatory subunit to the immunological synapse. *J Immunol* 2008;181:1969–77.
183. Nurieva RI, Chung Y, Hwang D, Yang XO, Kang HS, Ma L, et al. Generation of T follicular helper cells is mediated by interleukin-21 but independent of T helper 1, 2, or 17 cell lineages. *Immunity* 2008;29:138–49.
184. Choi YS, Kageyama R, Eto D, Escobar TC, Johnston RJ, Monticelli L, et al. ICOS receptor instructs T follicular helper cell versus effector cell differentiation via induction of the transcriptional repressor Bcl6. *Immunity* 2011;34:932–46.
185. Maazi H, Patel N, Sankaranarayanan I, Suzuki Y, Rigas D, Soroosh P, et al. ICOS:ICOS-ligand interaction is required for type 2 innate lymphoid cell function, homeostasis, and induction of airway hyperreactivity. *Immunity* 2015;42:538–51.
186. Leconte J, Bagherzadeh Yazdchi S, Panneton V, Suh WK. Inducible costimulator (ICOS) potentiates TCR-induced calcium flux by augmenting PLCgamma1 activation and actin remodeling. *Mol Immunol* 2016;79:38–46.
187. Paterson DJ, Jefferies WA, Green JR, Brandon MR, Corthesy P, Puklavec M, et al. Antigens of activated rat T lymphocytes including a molecule of 50,000 Mr detected only on CD4 positive T blasts. *Mol Immunol* 1987;24:1281–90.
188. Godfrey WR, Fagnoni FF, Harara MA, Buck D, Engleman EG. Identification of a human OX-40 ligand, a costimulator of CD4+ T cells with homology to tumor necrosis factor. *J Exp Med* 1994;180:757–62.
189. Gramaglia I, Weinberg AD, Lemon M, Croft M. OX-40 ligand: a potent costimulatory molecule for sustaining primary CD4 T cell responses. *J Immunol* 1998;161:6510–7.
190. Rogers PR, Song J, Gramaglia I, Killeen N, Croft M. OX40 promotes Bcl-xL and Bcl-2 expression and is essential for long-term survival of CD4 T cells. *Immunity* 2001;15:445–55.
191. Zingoni A, Sornasse T, Cocks BG, Tanaka Y, Santoni A, Lanier LL. Cross-talk between activated human NK cells and CD4+ T cells via OX40-OX40 ligand interactions. *J Immunol* 2004;173:3716–24.
192. Baumann R, Yousefi S, Simon D, Russmann S, Mueller C, Simon HU. Functional expression of CD134 by neutrophils. *Eur J Immunol* 2004;34:2268–75.
193. Song J, Salek-Ardakani S, Rogers PR, Cheng M, Van Parijs L, Croft M. The costimulation-regulated duration of PKB activation controls T cell longevity. *Nat Immunol* 2004;5:150–8.
194. Nocentini G, Giunchi L, Ronchetti S, Krausz LT, Bartoli A, Moraca R, et al. A new member of the tumor necrosis factor/nerve growth factor receptor family inhibits T cell receptor-induced apoptosis. *Proc Natl Acad Sci U S A* 1997;94:6216–21.
195. Gurney AL, Marsters SA, Huang RM, Pitti RM, Mark DT, Baldwin DT, et al. Identification of a new member of the tumor necrosis factor family and its receptor, a human ortholog of mouse GITR. *Curr Biol* 1999;9:215–8.
196. Kwon B, Yu KY, Ni J, Yu GL, Jang IK, Kim YJ, et al. Identification of a novel activation-inducible protein of the tumor necrosis factor receptor superfamily and its ligand. *J Biol Chem* 1999;274:6056–61.
197. McHugh RS, Whitters MJ, Piccirillo CA, Young DA, Shevach EM, Collins M, et al. CD4(+)CD25(+) immunoregulatory T cells: gene expression analysis reveals a functional role for the glucocorticoid-induced TNF receptor. *Immunity* 2002;16:311–23.
198. Shimizu J, Yamazaki S, Takahashi T, Ishida Y, Sakaguchi S. Stimulation of CD25(+)CD4(+) regulatory T cells through GITR breaks immunological self-tolerance. *Nat Immunol* 2002;3:135–42.
199. Hanabuchi S, Watanabe N, Wang YH, Wang YH, Ito T, Shaw J, et al. Human plasmacytoid dendritic cells activate NK cells through glucocorticoid-induced tumor necrosis factor receptor-ligand (GITRL). *Blood* 2006;107:3617–23.
200. Esparza EM, Lindsten T, Stockhausen JM, Arch RH. Tumor necrosis factor receptor (TNFR)-associated factor 5 is a critical intermediate of costimulatory signaling pathways triggered by glucocorticoid-induced TNFR in T cells. *J Biol Chem* 2006;281:8559–64.
201. Kwon BS, Weissman SM. cDNA sequences of two inducible T-cell genes. *Proc Natl Acad Sci U S A* 1989;86:1963–7.
202. Pollok KE, Kim YJ, Zhou Z, Hurtado J, Kim KK, Pickard RT, et al. Inducible T cell antigen 4-1BB. Analysis of expression and function. *J Immunol* 1993;150:771–81.
203. Saoulli K, Lee SY, Cannons JL, Yeh WC, Santana A, Goldstein MD, et al. CD28-independent, TRAF2-dependent costimulation of resting T cells by 4-1BB ligand. *J Exp Med* 1998;187:1849–62.
204. Futagawa T, Akiba H, Kodama T, Takeda K, Hosoda Y, Yagita H, et al. Expression and function of 4-1BB and 4-1BB ligand on murine dendritic cells. *Int Immunol* 2002;14:275–86.
205. Wilcox RA, Chapoval AI, Gorski KS, Otsuji M, Shin T, Flies DB, et al. Cutting edge: expression of functional CD137 receptor by dendritic cells. *J Immunol* 2002;168:4262–7.
206. Banchereau J, Dubois B, Fayette J, Burdin N, Brière F, Miossec P, et al. Functional CD40 antigen on B cells, dendritic cells and fibroblasts. *Adv Exp Med Biol* 1995;378:79–83.
207. Garside P, Ingulli E, Merica RR, Johnson JG, Noelle RJ, Jenkins MK. Visualization of specific B and T lymphocyte interactions in the lymph node. *Science* 1998;281:96–9.
208. Ahonen C, Manning E, Erickson LD, O'Connor B, Lind EF, Pullen SS, et al. The CD40-TRAF6 axis controls affinity maturation and the generation of long-lived plasma cells. *Nat Immunol* 2002;3:451–6.
209. Mackey MF, Wang Z, Eichelberg K, Germain RN. Distinct contributions of different CD40 TRAF binding sites to CD154-induced dendritic cell maturation and IL-12 secretion. *Eur J Immunol* 2003;33:779–89.
210. Goodwin RG, Alderson MR, Smith CA, Armitage RJ, VandenBos T, Jerzy R, et al. Molecular and biological characterization of a ligand for CD27 defines a new family of cytokines with homology to tumor necrosis factor. *Cell* 1993;73:447–56.
211. Gravestien LA, Blom B, Nolten LA, de Vries E, van der Horst G, Ossendorp F, et al. Cloning and expression of murine CD27: comparison with 4-1BB, another lymphocyte-specific member of the nerve growth factor receptor family. *Eur J Immunol* 1993;23:943–50.
212. Akiba H, Nakano H, Nishinaka S, Shindo M, Kobata T, Atsuta M, et al. CD27, a member of the tumor necrosis factor receptor superfamily, activates NF-kappaB and stress-activated protein kinase/c-Jun N-terminal kinase via TRAF2, TRAF5, and NF-kappaB-inducing kinase. *J Biol Chem* 1998;273:13353–8.
213. Hendriks J, Gravestien LA, Tesselaar K, van Lier RA, Schumacher TN, Borst J. CD27 is required for generation and long-term maintenance of T cell immunity. *Nat Immunol* 2000;1:433–40.
214. Hayakawa Y, Smyth MJ. CD27 dissects mature NK cells into two subsets with distinct responsiveness and migratory capacity. *J Immunol* 2006;176:1517–24.

CANCER DISCOVERY

Fundamental Mechanisms of Immune Checkpoint Blockade Therapy

Spencer C. Wei, Colm R. Duffy and James P. Allison

Cancer Discov 2018;8:1069-1086. Published OnlineFirst August 16, 2018.

Updated version Access the most recent version of this article at:
doi:[10.1158/2159-8290.CD-18-0367](https://doi.org/10.1158/2159-8290.CD-18-0367)

Cited articles This article cites 214 articles, 99 of which you can access for free at:
<http://cancerdiscovery.aacrjournals.org/content/8/9/1069.full#ref-list-1>

Citing articles This article has been cited by 2 HighWire-hosted articles. Access the articles at:
<http://cancerdiscovery.aacrjournals.org/content/8/9/1069.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cancerdiscovery.aacrjournals.org/content/8/9/1069>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.