

## Exercise One

1. Show that  $\int_0^t \int_0^{t_1} \cdots \int_0^{t_{k-1}} dt_k \cdots dt_1 = \frac{t^k}{k!}$ .
2. Prove Theorem 2.3, Taylor expansion, multi-index version.
3. Prove Theorem 2.8, tensor as multilinear maps
4. Recall that a function  $f : [a, b] \rightarrow \mathbb{R}$  is absolutely continuous if for every  $\varepsilon > 0$ , there exists a  $\delta > 0$  such that

$$\sum_{i=1}^n |a_i - a_{i+1}| < \delta \implies \sum_{i=1}^n |f(a_i) - f(a_{i+1})| < \varepsilon,$$

whenever  $[a_1, a_2], \dots, [a_n, a_{n+1}]$  are nonoverlapping subintervals of  $[a, b]$ .

- Check the construction of the Cantor function  $c : [0, 1] \rightarrow [0, 1]$
- Is the Cantor function absolutely continuous?
- Show that if  $f : [a, b] \rightarrow \mathbb{R}$  is absolutely continuous, and if  $\varphi : [a, b] \rightarrow \mathbb{R}$  is a smooth function that vanishes at the end points of the interval, then

$$(0.1) \quad \int_a^b f'(x)\varphi(x)dx = - \int_a^b f(x)\varphi'(x)dx.$$

5\*. Try whether you can prove the following statement: if  $f : [a, b] \rightarrow \mathbb{R}$  is an integrable function such that

$$(0.2) \quad \int_a^b g(x)\varphi(x)dx = - \int_a^b f(x)\varphi'(x)dx$$

for some integrable  $g : [a, b] \rightarrow \mathbb{R}$  and for all smooth function  $\varphi : [a, b] \rightarrow \mathbb{R}$  is a smooth function that vanishes at the end points of the interval, then  $f$  agrees almost everywhere with an absolutely continuous function; moreover,  $f' = g$  almost everywhere in this case.

Hint: You are able to use the following fact: Let  $f \in L^1(I)$  be such that

$$\int_I f\varphi' dx = 0$$

for all  $\varphi \in C_0^1(I)$ . Show that there exists a constant  $c$  such that  $f = c$  a.e. on  $I$ .