

Equilibria of collisionless systems

1rd part

Outlines

Weak bars

- the Lindblad resonances
- orbit families in realistic bars

The collisionless Boltzmann equation

- The distribution function (DF) of stellar systems
- The Collisionless Boltzmann equation
- Limitations

Relations between DFs and observables

- Density, velocity distribution function, mean velocity, velocity dispersion


The Jeans theorems

- Solutions of the Collisionless Boltzmann equation
- Symmetries and DFs

Stellar Orbits

Orbits in weak rotating bars

The weakly-bared galaxy model



General solution

(harmonic oscillator of frequ. ω_0 driven at frequ. $m(\Omega_0 - \Omega_1)$)

$$R_n(t) = C_n \cos(\omega_0 t + \alpha) - \left[\frac{d\phi_3}{dR} + \frac{2\Omega\phi_0}{R(\Omega - \Omega_1)} \right]_{R_0} \frac{\cos(m(\Omega_0 - \Omega_1)t)}{\omega_0^2 - m^2(\Omega_0 - \Omega_1)^2}$$

using $\varphi_0(t) = (\Omega_0 - \Omega_1)t$

$$R_n(\varphi_0) = C_n \cos\left(\frac{\omega_0 \varphi_0}{\Omega_0 - \Omega_1} + \alpha\right) - \left[\frac{d\phi_3}{dR} + \frac{2\Omega\phi_0}{R(\Omega - \Omega_1)} \right]_{R_0} \frac{\cos(m\varphi_0)}{\omega_0^2 - m^2(\Omega_0 - \Omega_1)^2}$$

C_n, α arbitrary constants

$$\dot{\varphi}_n = -2\Omega_0 \frac{R_n}{R_0} - \frac{\phi_3(R_0)}{\Omega_0^2 (\Omega_0 - \Omega_1)} \cos(m(\Omega_0 - \Omega_1)t) + cte$$

Discussion

$$R_1(\varphi_0) = C_1 \cos\left(\frac{x_0 \varphi_0}{R_0 - R_1} + \alpha\right) - \left[\frac{d\phi_3}{dR} + \frac{2\Omega\phi_3}{R(\Omega - \Omega_1)} \right]_{R_0} \frac{\cos(m\varphi_0)}{x_0^2 - m^2(R_0 - R_1)^2}$$

① if $\phi_3(R) = 0$ (no perturbation)

$$R_1(t) = C_1 \cos(x_0 t + \alpha)$$

$$\dot{\varphi}_1(t) = -2\Omega_0 \frac{R_1(t)}{R_0}$$

Epicycles motions

$\equiv x(t)$ radial oscillations

$\Rightarrow y(t)$ oscillations along the orbit

② if $C_1 = 0$ $\phi_3 \neq 0$

$$R_1(\varphi_0) = - \left[\frac{d\phi_3}{dR} + \frac{2\Omega\phi_3}{R(\Omega - \Omega_1)} \right]_{R_0} \frac{\cos(m\varphi_0)}{x_0^2 - m^2(R_0 - R_1)^2}$$

cte

periodic in φ_0 ($\frac{2\pi}{m}$)

\Rightarrow closed orbit



③ if $C_1 \neq 0$ oscillations around the closed orbit (same family)

The orbit is not necessary closed

Resonances



two problematic terms

$$\frac{1}{\Omega_0 - \Omega_b} \quad \text{and} \quad \frac{1}{\kappa^2 - m^2(\Omega_0 - \Omega_b)^2}$$

$\Rightarrow R_1$ may diverge !

1)

$$\Omega_0 = \Omega_b$$

Corotation

we are at a radius where the circular frequency is similar to the pattern speed of the bar

as $\dot{\varphi}_0 = \Omega_0 - \Omega_b \Rightarrow \underline{\dot{\varphi}_0 = 0}$

\rightarrow stable in the rotating frame

2)

$$m(\Omega_0 - \Omega_b) = \pm \kappa$$

Lindblad resonances

freq. at which the star encounter the potential minimum

\rightarrow the frequency at which a star encounter a potential minimum is similar to its radial frequency

$$\equiv \Omega_b = \Omega \pm \frac{\kappa}{m}$$


\Rightarrow excitation

A circular orbit has two natural frequencies

-
- ① κ : radial frequ. \rightarrow
 - ② 0 : azimuthal frequ. \rightarrow
(no change \Rightarrow frequ. = 0)

Resonances occur when the forcing frequency $m(\Omega_0 - \Omega_b)$ is equal to one of these frequencies.

Disk : Miyamoto-Nagai
 Bulge : Plummer



Inner Lindblad resonances
 (ILR1, ILR2)

$$\Omega_b = \Omega - \kappa/2$$

the orbit rotate faster than the
 bar



Corotation (CR)

$$\Omega_b = \Omega$$

Outer Lindblad resonance
 (OLR)



$$\Omega_b = \Omega + \kappa/2$$

the orbit rotate slower than the
 bar





$R = 0.2$

$R < R_{ILR1}$





$R = 0.3$

$R \cong R_{ILR1}$





$$R = 0.7$$

$$R_{\text{ILR1}} < R < R_{\text{ILR2}}$$





$R = 0.9$

$R \cong R_{ILR2}$





$$R = 1.5$$

$$R_{\text{ILR2}} < R < R_{\text{CR}}$$




$$R = 2.0$$

$$R_{\text{ILR2}} < R < R_{\text{CR}}$$





$$R = 3.0$$

$$R_{\text{ILR2}} < R < R_{\text{CR}}$$



$R = 3.5$



$R \cong R_{CR}$



+ closed orbits
around L4, L5



$R = 4.0$

$R_{\text{CR}} < R < R_{\text{OLR}}$





$$R = 4.5$$

$$R_{\text{CR}} < R < R_{\text{OLR}}$$





$R = 5.0$

$R_{\text{CR}} < R < R_{\text{OLR}}$





$R = 5.5$

$R \cong R_{\text{OLR}}$





$R = 6.0$

$R_{\text{OLR}} < R$




$R = 7.0$


$R_{\text{OLR}} < R$





$$R = 8.0$$

$$R_{\text{OLR}} < R$$








Bifurcation : apparition of x2 (stable)/x3 (unstable) orbits





```
./mapping.py --V0 1. --q 0.8 --Rc 0.03 --Omega 1 -E -2.5 --x 0.0268
```

x1 : prograde x4 : retrograde





```
./mapping.py --V0 1. --q 0.8 --Rc 0.03 --Omega 1 -E -2.5 --x 0.0766659  
./mapping.py --V0 1. --q 0.8 --Rc 0.03 --Omega 1 -E -2.5 --x -0.034
```

Lindblad frequencies for the Logarithmic potential




R_{ILR2}

R_{ILR1}



Buta & Combes 1998



Equilibria of collisionless systems

The collisionless Boltzmann equation

Introduction / Motivations

So far, we :

1. we modelled static potentials from a mass distribution (Poisson equation)
2. from the potential, we obtained forces and derived equations of motion leading us study orbits in different idealized potentials :
 - spherical potentials
 - axi-symmetric potentials (epicycles motions)
 - orbits in bared rotating potentials (motions around Lagrange points)

But :

1. We did not used any velocity constraints. We only used the positions of stars through the emission of light.
2. Nothing tells us that the models we used are at the equilibrium.
This is not guarantee, if, for e.g., all velocities are zero...
3. We did do not include the self-gravity of the model or perturbations on it due to the orbits of stars.

Introduction / Motivations

Goal :

Build a self-consistent way galaxies, ensuring that they are at the equilibrium, i.e., if we compute the evolution of the galaxy under its own gravity, the evolution will be stationary.

→ requires the description of the **density** but also the **velocity field**

$$\rho(\vec{x}) \qquad \vec{v}(\vec{x})$$

Assumptions :

1. We will consider systems with a very large number of “particles” (stars, DM)

→ the collisionless approximation is valid

→ real orbits deviates not too much from the one predicted from the model
(very large relaxation time)

We will seek for solution corresponding to $t_{\text{relax}} = \infty$

2. We will consider systems composed of N identical particles, i.e., with all the same mass.

All particles will be equivalent

Introduction / Motivations

Goal :

Build a self-consistent way galaxies, ensuring that they are at the equilibrium, i.e., if we compute the evolution of the galaxy under its own gravity, the evolution will be stationary.

→ requires the description of the **density** but also the **velocity field**

$$\rho(\vec{x})$$

$$\vec{v}(\vec{x})$$

But :

It is impossible to describe analytically the orbits of billions of stars :

→ **we need a probabilistic approach**

Distribution function (DF)

Definition ① $f(\vec{x}, \vec{v}, t)$ or $f(\vec{w}, t)$ such that
 $f(\vec{x}, \vec{v}, t) d^3\vec{x} d^3\vec{v}$ or $f(\vec{w}, t) d^3\vec{w}$
is the probability that at the time t ,
a randomly chosen star "i" has its position \vec{x}_i ,
an velocity \vec{v}_i , or phase space coordinates \vec{w}_i
in the ranges $\vec{x}_i \in [\vec{x}, \vec{x} + d\vec{x}]$
 $\vec{v}_i \in [\vec{v}, \vec{v} + d\vec{v}]$
 $\equiv \vec{w}_i \in [\vec{w}, \vec{w} + d\vec{w}]$

obviously:
(normalisation)

$$\int f(\vec{x}, \vec{v}, t) d^3\vec{x} d^3\vec{v} = 1$$
$$\equiv \int f(\vec{w}, t) d^3\vec{w} = 1$$

the particle
is for sure
somewhere in
the phase space

$f(\vec{x}, \vec{v}, t)$ is the probability density of the phase space.

Distribution function (DF)

Definition (2) $\tilde{f}(\vec{x}, \vec{v}, t)$ or $\tilde{f}(\vec{w}, t)$ such that

$$\tilde{f}(\vec{x}, \vec{v}, t) d^3\vec{x} d^3\vec{v} \quad \text{or} \quad \tilde{f}(\vec{w}, t) d^3\vec{w}$$

is the number of stars having position \vec{x} and velocities \vec{v} (\vec{w}) in the intervals at time t :

$$\vec{x}_i \in [\vec{x}, \vec{x} + d\vec{x}]$$

$$\vec{v}_i \in [\vec{v}, \vec{v} + d\vec{v}]$$

$$\vec{w}_i \in [\vec{w}, \vec{w} + d\vec{w}]$$

obviously:

(normalisation)

$$\int \tilde{f}(\vec{x}, \vec{v}, t) d^3\vec{x} d^3\vec{v} = N$$

$$\equiv \int \tilde{f}(\vec{w}, t) d^6\vec{w} = N$$

There are exactly N particles in the phase space

$\tilde{f}(\vec{x}, \vec{v}, t)$ is the number density of the phase space.

Combining Det. ① and Det ②

$$N f(\bar{x}, \bar{v}, t) = \tilde{f}(\bar{x}, \bar{v}, t)$$

Notes

- we will sometimes forget the " \sim "
- the time dependence " t " will not be systematically written

Using definition ①

The probability of finding a star "i" in the subvolume of the phase space γ is:

$$P = \int_{\gamma} f(\vec{w}) d^6\vec{w}$$

However, imagine that we are using another canonical coordinate system \vec{W} (in which the Hamilton equations are valid)

e.g. $(x, y, p_x = \dot{x}, p_y = \dot{y}) \rightarrow (r, \theta, p_r = \dot{r}, p_\theta = r^2\dot{\theta})$

$$P^w = \int_{\gamma} F(\vec{W}) d^6\vec{W} = P$$

The probability must not be affected by a coordinate change.

If ν is taken small enough, we can assume $g(\vec{w})$ and $F(\vec{W})$ to be constant and hence

$$g(\vec{w}_\nu) \int_\nu d^6\vec{w} = F(\vec{W}_\nu) \int_\nu d^6\vec{W}$$

But, for canonical coordinates, the phase space volume element is the same:

$$\int_\nu d^6\vec{w} = \int_\nu d^6\vec{W}$$

Thus

$$g(\vec{w}) = F(\vec{W})$$

The density of the phase space is independent of the coordinate system

Corollary: We can use any system of canonical coordinates $\vec{w} = (\vec{q}, \vec{p})$ to define the distribution function

The collisionless Boltzmann equation

- What is the evolution of $f(\vec{w})$ over time?

As the mass, the probability is a conserved quantity. $\rho = N f$

the number of stars is a conserved quantity.

in the phase space

Continuity equation (similar than for hydrodynamics)



the time variation of the mass in V

$$\frac{dM}{dt} = \sum_{\text{faces}} \rho \vec{v} \cdot d\vec{S}$$

mass flux

Mass conservation

$$\frac{\partial \rho}{\partial t} + \vec{\nabla}_x \cdot (\rho \vec{v}) = 0$$

Probability conservation

$$\frac{\partial f}{\partial t} + \vec{\nabla}_w \cdot (f \vec{w}) = 0$$

mass flux through the surface
of the volume

probability flux through the surface
of the volume

Analogy with the continuity equation in hydrodynamics

$$f(\vec{x}, t) \quad \vec{v} = \frac{d}{dt} \vec{x}$$

$$\frac{\partial f}{\partial t} + \vec{\nabla}_{\vec{x}} \cdot (f \vec{v}) = 0$$

Lagrangian derivative

$$\frac{d}{dt} f(\vec{x}, t) = \frac{\partial f}{\partial t} + \vec{v} \cdot \vec{\nabla}_{\vec{x}} f$$

Flux divergence

$$\vec{\nabla}_{\vec{x}} \cdot (f \vec{v}) = \vec{v} \cdot \vec{\nabla}_{\vec{x}} f + f \vec{\nabla}_{\vec{x}} \cdot \vec{v}$$

$$\vec{v} \cdot \vec{\nabla}_{\vec{x}} f = \vec{\nabla}_{\vec{x}} \cdot (f \vec{v}) - f \vec{\nabla}_{\vec{x}} \cdot \vec{v}$$

$$f(\vec{w}, t) \quad \dot{\vec{w}} = \frac{d}{dt} \vec{w}$$

$$\frac{\partial f}{\partial t} + \vec{\nabla}_{\vec{w}} \cdot (f \dot{\vec{w}}) = 0$$

Lagrangian derivative

$$\frac{d}{dt} f(\vec{w}, t) = \frac{\partial f}{\partial t} + \dot{\vec{w}} \cdot \vec{\nabla}_{\vec{w}} f$$

Flux divergence

$$\vec{\nabla}_{\vec{w}} \cdot (f \dot{\vec{w}}) = \dot{\vec{w}} \cdot \vec{\nabla}_{\vec{w}} f + f \vec{\nabla}_{\vec{w}} \cdot \dot{\vec{w}}$$

$$\dot{\vec{w}} \cdot \vec{\nabla}_{\vec{w}} f = \vec{\nabla}_{\vec{w}} \cdot (f \dot{\vec{w}}) - f \vec{\nabla}_{\vec{w}} \cdot \dot{\vec{w}}$$

Lagrangian derivative

$$\begin{aligned}\frac{d}{dt} f(\vec{x}, t) &= \frac{\partial f}{\partial t} + \vec{v} \cdot \vec{\nabla}_x f \\ &= \underbrace{\frac{\partial f}{\partial t} + \vec{\nabla}_x (f \vec{v})}_{= 0 \text{ continuity Eq.}} - \rho \vec{\nabla}_x \cdot \vec{v}\end{aligned}$$

$$\frac{d}{dt} f(\vec{x}, t) = - \rho \vec{\nabla}_x \cdot \vec{v}$$

the increase of f along the flow is due to compression

incompressible fluid :

$$\vec{\nabla}_x \cdot \vec{v} = 0$$

Lagrangian derivative

$$\begin{aligned}\frac{d}{dt} f(\vec{w}, t) &= \frac{\partial f}{\partial t} + \dot{\vec{w}} \cdot \vec{\nabla}_w f \\ &= \underbrace{\frac{\partial f}{\partial t} + \vec{\nabla}_w (f \dot{\vec{w}})}_{= 0 \text{ continuity Eq.}} - \underbrace{\rho \vec{\nabla}_w \cdot \dot{\vec{w}}}_{= 0 \text{ canonical coords.}}\end{aligned}$$

$$\frac{d}{dt} f(\vec{w}, t) = 0$$

\Rightarrow behaves like an incompressible fluid

The flow through the phase space is incompressible

Seen from an observer that follows the flow in the phase space, i.e. an orbit : f is constant

Expressing the continuity equation using $\tilde{\omega} = (\tilde{q}, \tilde{p})$

$$\frac{d}{dt} f(\tilde{\omega}, t) = \frac{\partial f(\tilde{\omega}, t)}{\partial t} + \vec{\nabla}_{\tilde{\omega}} (f(\tilde{\omega}, t) \dot{\tilde{\omega}}) = 0$$

$$= \frac{\partial f(\tilde{\omega}, t)}{\partial t} + \dot{\tilde{\omega}} \cdot \vec{\nabla}_{\tilde{\omega}} (f(\tilde{\omega}, t)) = 0$$

$$= \frac{\partial f(\tilde{q}, \tilde{p})}{\partial t} + \sum_i \dot{q}_i \frac{\partial}{\partial q_i} f(\tilde{q}, \tilde{p}) + \sum_i \dot{p}_i \frac{\partial}{\partial p_i} f(\tilde{q}, \tilde{p})$$

$$\frac{d}{dt} f(\tilde{\omega}, t) = \frac{\partial f(\tilde{q}, \tilde{p})}{\partial t} + \dot{\tilde{q}} \cdot \frac{\partial}{\partial \tilde{q}} f(\tilde{q}, \tilde{p}) + \dot{\tilde{p}} \cdot \frac{\partial}{\partial \tilde{p}} f(\tilde{q}, \tilde{p}) = 0$$

The Collisionless Boltzmann Equation

Using the Hamilton Equations

$$\dot{\vec{q}} = \frac{\partial H}{\partial \vec{p}} \quad \dot{\vec{p}} = - \frac{\partial H}{\partial \vec{q}}$$

Then

$$\frac{\partial}{\partial t} f + \dot{\vec{q}} \frac{\partial}{\partial \vec{q}} f + \dot{\vec{p}} \frac{\partial}{\partial \vec{p}} f = 0$$

becomes

$$\frac{\partial}{\partial t} f + \frac{\partial H}{\partial \vec{p}} \frac{\partial f}{\partial \vec{q}} - \frac{\partial H}{\partial \vec{q}} \frac{\partial f}{\partial \vec{p}} = 0$$

$$\frac{\partial}{\partial t} f + [f, H] = 0$$

Poisson brackets

$$[A, B] := \frac{\partial A}{\partial \vec{q}} \frac{\partial B}{\partial \vec{p}} - \frac{\partial A}{\partial \vec{p}} \frac{\partial B}{\partial \vec{q}}$$
$$= \sum_i^n \frac{\partial A}{\partial q_i} \frac{\partial B}{\partial p_i} - \frac{\partial A}{\partial p_i} \frac{\partial B}{\partial q_i}$$

Liouville's Theorem (corollary)

In the motion of a stellar system, any volume of phase space remains constant


dV : an infinitesimal volume of the phase space

$dN(t)$: the number of stars in $dV(t)$ at t

$$dN(t) = \tilde{f}(\tilde{w}, t) dV(t)$$

$dN(t')$: the number of stars in $dV(t')$ at t'

$$dN(t') = \tilde{f}(\tilde{w}, t') dV(t')$$



$$\text{But } dN(t) = dN(t')$$

$$\equiv \frac{dN}{dt} = 0$$

Because EoM are 1st order differential equations, only the points that were in dV at t are in dV' at t'

Thus

$$\frac{dN}{dt} = \frac{d}{dt} \left(\tilde{f}(w, t) dV(t) \right)$$


$$= \underbrace{\frac{d}{dt} \left(\tilde{f}(w, t) \right)}_{=0 \text{ (Boltzmann equation)}} dV(t) + \tilde{f}(w, t) \frac{d}{dt} (dV(t)) = 0$$


$$\Rightarrow \frac{d}{dt} (dV(t))$$

$$dV(t) = \text{cte}$$

The distribution function remains constant along the flow


Illustration 1 : Ideal race: each runner has a constant speed





The distribution function remains constant along the flow

Illustration 1 : Ideal race: each runner has a constant speed



ν : the phase space volume

$$\tilde{f}(t = 0) = \frac{N}{\nu_0} = \frac{N}{\Delta x \Delta v}$$

$$\tilde{f}(t = t) = \frac{N}{\nu_t} = \frac{N}{\Delta x \Delta v}$$

Illustration 2 : Harmonic oscillator

$$H(x, \dot{x}) = \frac{1}{2}\dot{x}^2 + \frac{1}{2}\omega x^2$$

$$\omega = 1$$

2.0


$$\begin{cases} x(t) = A \cos(\omega t + \phi_0) \\ v(t) = -A\omega \sin(\omega t + \phi_0) \end{cases}$$

$$\begin{cases} x^2(t) = A^2 \cos^2(\omega t + \phi_0) \\ v^2(t) = A^2\omega^2 \sin^2(\omega t + \phi_0) \end{cases}$$

$$cte^2 = x^2 + \frac{v^2}{\omega^2}$$

particles move along ellipses in the phase space

For : $\omega = 1$ pure rotation



-2.0

-2.0 -1.5 -1.0 -0.5 0.0 0.5 1.0 1.5 2.0

x






Illustration 3 : Harmonic oscillator

$$H(x, \dot{x}) = \frac{1}{2}\dot{x}^2 + \frac{1}{2}\omega x^2$$

$$\omega = 0.75$$





The Collisionless Boltzmann equation in various coordinates

Generalized coordinates

$$\vec{p} = \frac{\partial L(\vec{q}, \dot{\vec{q}})}{\partial \dot{\vec{q}}}$$

$$\frac{\partial f}{\partial t} + \dot{\vec{q}} \cdot \frac{\partial f}{\partial \vec{q}} + \dot{\vec{p}} \cdot \frac{\partial f}{\partial \vec{p}} \equiv \frac{\partial f}{\partial t} + \frac{\partial f}{\partial \vec{q}} \cdot \frac{\partial H}{\partial \vec{p}} - \frac{\partial f}{\partial \vec{p}} \cdot \frac{\partial H}{\partial \vec{q}} = 0$$

Cartesian coordinates

$$\begin{cases} p_x = \dot{x} = v_x \\ p_y = \dot{y} = v_y \\ p_z = \dot{z} = v_z \end{cases} \quad H = \frac{1}{2} (v_x^2 + v_y^2 + v_z^2) + \Phi(x, y, z)$$

$$\frac{\partial f}{\partial t} + \vec{v} \cdot \frac{\partial f}{\partial \vec{x}} - \frac{\partial \Phi}{\partial \vec{x}} \cdot \frac{\partial f}{\partial \vec{v}} = 0$$

Spherical coordinates

$$\begin{cases} p_r = \dot{r} = v_r \\ p_\theta = r^2 \dot{\theta} = r v_\theta \\ p_\phi = r^2 \sin^2(\theta) \dot{\phi} = r \sin(\theta) v_\phi \end{cases}$$

$$H = \frac{1}{2} \left(p_r^2 + \frac{p_\theta^2}{r^2} + \frac{p_\phi^2}{r^2 \sin^2(\theta)} \right) + \Phi(R, \theta, \phi)$$

$$\frac{\partial f}{\partial t} + p_r \frac{\partial f}{\partial r} + \frac{p_\theta}{r^2} \frac{\partial f}{\partial \theta} + \frac{p_\phi}{r^2 \sin^2(\theta)} \frac{\partial f}{\partial \phi} - \left(\frac{\partial \Phi}{\partial r} - \frac{p_\theta^2}{r^3} - \frac{p_\phi^2}{r^3 \sin^2(\theta)} \right) \frac{\partial f}{\partial p_r} - \left(\frac{\partial \Phi}{\partial \theta} - \frac{p_\phi^2 \cos(\theta)}{r^2 \sin^3(\theta)} \right) \frac{\partial f}{\partial p_\theta} - \frac{\partial \Phi}{\partial \phi} \frac{\partial f}{\partial p_\phi} = 0$$

Cylindrical coordinates

$$\begin{cases} p_R = \dot{R} = v_R \\ p_\phi = R^2 \dot{\phi} = R v_\phi \\ p_z = \dot{z} = v_z \end{cases}$$

$$H = \frac{1}{2} \left(p_R^2 + \frac{p_\phi^2}{R^2} + p_z^2 \right) + \Phi(R, \phi, z)$$

$$\frac{\partial f}{\partial t} + p_R \frac{\partial f}{\partial R} + \frac{p_\phi}{R^2} \frac{\partial f}{\partial \phi} + p_z \frac{\partial f}{\partial z} - \left(\frac{\partial \Phi}{\partial R} - \frac{p_\phi^2}{R^3} \right) \frac{\partial f}{\partial p_R} - \frac{\partial \Phi}{\partial \phi} \frac{\partial f}{\partial p_\phi} - \frac{\partial \Phi}{\partial z} \frac{\partial f}{\partial p_z} = 0$$

Limits of the Collisionless Boltzmann equation

I. Finite stellar lifetime

- Stars are created and die. The hypothesis of conservation of the probability/number is violated.

We should better have (in Cartesian coordinates):

$$\frac{\partial f}{\partial t} + \vec{v} \cdot \frac{\partial f}{\partial \vec{x}} - \frac{\partial \Phi}{\partial \vec{x}} \cdot \frac{\partial f}{\partial \vec{v}} = B(\vec{x}, \vec{v}, t) - D(\vec{x}, \vec{v}, t)$$

$\sim \frac{v}{R} f$
 $\sim \frac{a}{v} f$

$\sim \frac{1}{t_{\text{cross}}} f$
 $\sim \frac{1}{t_{\text{cross}}} f$

Rate per unit phase-space
volume at which stars are
born and die

- Define

$$\gamma = \frac{|B - D|}{f} t_{\text{cross}}$$

If $\gamma \ll 1$ the approximation is ok

i.e. : the fractional change in the number of stars per crossing time must be small.

Limits of the Collisionless Boltzmann equation

Examples:

- M-stars in an elliptical galaxies
 - Life time > 10 Gyr ($> t_{\text{cross}}$) $\gamma \cong 0$
 - $B=0$ (no star formation)
- O-stars in the Milky Way
 - Life time < 100 Myr ($< t_{\text{cross}}$) $\gamma \gg 1$
 - Do not move much, the phase space distribution will be dominated by star formation processes
- Main sequence stars ($M < 1.5M_{\odot}$)
 - Life time > 1 Gyr ($> t_{\text{cross}}$) $\gamma \cong 0$

Limits of the Collisionless Boltzmann equation

II. Correlation between stars

- We assumed that the probability of finding one peculiar stars somewhere in the phase space is independent of the others. Mathematically: the probability of finding particle "i" in $d^6\vec{\omega}$ and "j" in $d^6\vec{\omega}'$ is :

$$f(\vec{\omega})d^6\vec{\omega} \cdot f(\vec{\omega}')d^6\vec{\omega}'$$

This is not completely true, as stars interact gravitationally and my generate correlations.

However, this is not a real problem as long as the forces between nearby stars do not dominates over the forces due to the rest of the system (the definition of a collisionless system).

Equilibria of collisionless systems

**Relations between the DFs
and observables**

Relations between the DF and observables

$$f(\vec{w})$$

- $f(\vec{w})$: probability density
in the phase space
- $f(\vec{w}) d^6\vec{w}$: probability of finding 1 star
in the phase space volume $[\vec{w}, \vec{w} + d\vec{w}]$

Distribution function in the configuration space

$$\nu(\vec{x}) = \int d^3\vec{v} f(\vec{x}, \vec{v})$$

- $\nu(\vec{x})$: probability density
in the configuration space
- $\nu(\vec{x}) d^3\vec{x}$: probability of finding 1 star
in the configuration space volume $[\vec{x}, \vec{x} + d\vec{x}]$

Distribution function in the configuration space

$$n(\vec{x}) = N \nu(\vec{x}) = \int d^3\vec{v} \tilde{f}(\vec{x}, \vec{v})$$

- $n(\vec{x})$: number density of star in the configuration space
- $n(\vec{x}) d^3\vec{x}$: probability of finding N stars in the configuration space volume $[\vec{x}, \vec{x} + d\vec{x}]$

Distribution function in the configuration space

$$\rho(\vec{x}) = N \cdot m \cdot \nu(\vec{x}) = m \int d^3\vec{v} \tilde{f}(\vec{x}, \vec{v})$$

m : mass of particles

- $\rho(\vec{x})$: mass density of star in the configuration space
- $\rho(\vec{x}) d^3\vec{x}$: probability of finding a mass $M = N \cdot m$ in the configuration space volume $[\vec{x}, \vec{x} + d\vec{x}]$


Distribution function in the velocity space

$$P_{\vec{x}}(\vec{v}) = \frac{f(\vec{x}, \vec{v})}{v(\vec{x})}$$

$$\int P_{\vec{x}}(\vec{v}) d^3\vec{v} = \frac{1}{v(\vec{x})} \int \underbrace{f(\vec{x}, \vec{v})}_{:= v(\vec{x})} d^3\vec{v} = 1$$

\equiv velocity distribution function (VDF)

- $P_{\vec{x}}(\vec{v})$: probability density at the position \vec{x} in the velocity space
- $P_{\vec{x}}(\vec{v}) d^3\vec{v}$: probability of finding 1 star in \vec{x} in the velocity space volume $[\vec{v}, \vec{v} + d\vec{v}]$




can be measured near the sun

Mean velocity (first moment of the VDF)

$$\vec{V}(\vec{x}) = \int \vec{v} P_{\vec{x}}(\vec{v}) d^3\vec{v} = \frac{1}{V(\vec{x})} \int \vec{v} f(\vec{x}, \vec{v}) d^3\vec{v}$$

- along one peculiar axis \vec{n}



$$\vec{V}_{\vec{n}}(\vec{x}) = \int \vec{v} \cdot \vec{n} P_{\vec{x}}(\vec{v}) d^3\vec{v} = \frac{1}{V(\vec{x})} \int \vec{v} \cdot \vec{n} f(\vec{x}, \vec{v}) d^3\vec{v}$$

- if $\vec{n} = \vec{e}_i$


$$\vec{V}_i(\vec{x}) = \int v_i P_{\vec{x}}(\vec{v}) d^3\vec{v} = \frac{1}{V(\vec{x})} \int v_i f(\vec{x}, \vec{v}) d^3\vec{v}$$

Velocity dispersion tensor (second moment of the VDF)

$$\begin{aligned}\sigma_{ij}^2 &= \int (v_i - \bar{v}_i)(v_j - \bar{v}_j) P_{\vec{x}}(\vec{v}) d^3\vec{v} \\ &= \frac{1}{N(\vec{x})} \int (v_i - \bar{v}_i)(v_j - \bar{v}_j) f(\vec{x}, \vec{v}) d^3\vec{v} \\ &= \int v_i v_j f(\vec{x}, \vec{v}) d^3\vec{v} - \left(\int v_i f(\vec{x}, \vec{v}) d^3\vec{v} \right) \left(\int v_j f(\vec{x}, \vec{v}) d^3\vec{v} \right) \\ &= \overline{v_i v_j} - \bar{v}_i \bar{v}_j\end{aligned}$$

3x3 symmetric tensor
⇒ may be diagonalised


Describe an ellipsoid (velocity ellipsoid)



$$\sigma_{ij}^2 = \sigma_{ii}^2 \delta_{ij} = \begin{pmatrix} \sigma_{11} & 0 & 0 \\ 0 & \sigma_{22} & 0 \\ 0 & 0 & \sigma_{33} \end{pmatrix}$$

Illustration

$i = j$, 1D (x, v)



$$\sigma_{ii}^2(x) = \overline{v_i^2(x)} - \overline{v_i(x)}^2$$

"intrinsic"
dispersion

"total"
dispersion

"mean"

Equilibria of collisionless systems

The Jeans Theorems

Jeans theorems

Question : How can we solve the collisionless Boltzmann equation ?

Start with a 1-D steady state solution $\frac{\partial f}{\partial t} = 0$

$$\underbrace{\frac{\partial H}{\partial p}}_q \frac{\partial f}{\partial q} - \frac{\partial H}{\partial q} \frac{\partial f}{\partial p} = 0$$

In cartesian coordinates

$$\frac{\partial H}{\partial x} = \frac{\partial \phi}{\partial x}$$

$$\frac{\partial f}{\partial x} v - \frac{\partial \phi}{\partial x} \frac{\partial f}{\partial v} = 0$$

$$\frac{\partial f}{\partial x} v - \frac{\partial \phi}{\partial x} \frac{\partial f}{\partial v} = 0$$

linear homogeneous
differential equation

Solutions from Lagrange's equations

$$\frac{dx}{v} = - \frac{dV}{\frac{\partial \phi}{\partial x}}$$

$$- \frac{\partial \phi}{\partial x} dx = v dv$$

$$I(x, v) = \frac{1}{2} v^2 + \phi(x) \quad \text{is a solution}$$

$$dI(x, v) = \underbrace{v}_{\frac{\partial I}{\partial v}} dv + \underbrace{\frac{\partial \phi}{\partial x}}_{\frac{\partial I}{\partial x}} dx$$

The Hamiltonian $H(x, v)$ is a solution of the static CBE !

Back to the integrals of motion

The function $I(\tilde{x}(t), \tilde{v}(t))$ is an integral of motion if

$$\frac{d}{dt} I(\tilde{x}(t), \tilde{v}(t)) = 0$$

along the trajectory.

But

$$\frac{dI}{dt} = \frac{\partial I}{\partial \tilde{x}} \tilde{x}^i + \frac{\partial I}{\partial \tilde{v}} \tilde{v}^i = 0$$

$$= \frac{\partial I}{\partial \tilde{x}} \tilde{v} - \frac{\partial I}{\partial \tilde{v}} \tilde{v} \phi = 0$$

Similar to the
Collisionless Boltzmann
equation

If $I(\tilde{x}, \tilde{v})$ is an integral of motion

$I(\tilde{x}, \tilde{v})$ is a steady state solution of the

Collisionless Boltzmann equation




Jeans theorems

- I. Any steady-state solution of the collisionless Boltzmann equation depends on the phase-space coordinates only through integrals of motion in the given potential.

Demonstration:

If a function is a solution of the steady-state collisionless Boltzmann equation, then, it is an integral of motion, thus the function depends on the phase-space coordinates only through integrals of motion (itself !).

- II. Any function of integrals of motion yields a steady-state solution of the collisionless Boltzmann equation.



Jeans theorems

- I. Any steady-state solution of the collisionless Boltzmann equation depends on the phase-space coordinates only through integrals of motion in the given potential.


Demonstration:

If a function is a solution of the steady-state collisionless Boltzmann equation, then, it is an integral of motion, thus the function depends on the phase-space coordinates only through integrals of motion (itself !).

- II. Any function of integrals of motion yields a steady-state solution of the collisionless Boltzmann equation.

Demonstration:

Assume $f(\vec{x}, \vec{v}) = f(I_1(\vec{x}, \vec{v}), I_2(\vec{x}, \vec{v}), I_3(\vec{x}, \vec{v}), \dots)$ and derivate...



Jeans theorems

- I. Any steady-state solution of the collisionless Boltzmann equation depends on the phase-space coordinates only through integrals of motion in the given potential.

Demonstration:

If a function is a solution of the steady-state collisionless Boltzmann equation, then, it is an integral of motion, thus the function depends on the phase-space coordinates only through integrals of motion (itself !).

- II. Any function of integrals of motion yields a steady-state solution of the collisionless Boltzmann equation.

Extremely useful to generate DFs

Demonstration:

Assume $f(\vec{x}, \vec{v}) = f(I_1(\vec{x}, \vec{v}), I_2(\vec{x}, \vec{v}), I_3(\vec{x}, \vec{v}), \dots)$ and derivate...

Choices of DFs and relations with the velocity moments

1. DFs that depend only on H

(no particular symmetry)
except time!

Ergodic distribution functions

$$\phi = \phi(\vec{x}, t)$$

Example $\left\{ \begin{array}{l} H(\vec{x}, \vec{v}) = \frac{1}{2} \vec{v}^2 + \phi(\vec{x}) \\ f = f\left(\frac{1}{2} \vec{v}^2 + \phi(\vec{x})\right) \end{array} \right.$

Mean velocity

Note: the velocity dependency is only through v^2 (isotropic)

$$\vec{v}(\vec{x}) = \frac{1}{V(\vec{x})} \int \vec{v} f\left(\frac{1}{2} \vec{v}^2 + \phi(\vec{x})\right) d^3\vec{v} = 0$$

indeed

$$\bar{v}_x(\vec{x}) = \frac{1}{V(\vec{x})} \int_{-\infty}^{\infty} dv_z \int_{-\infty}^{\infty} dv_y \int_{-\infty}^{\infty} dv_x \underbrace{v_x}_{\text{odd}} \underbrace{f\left(\frac{1}{2} \vec{v}^2 + \phi(\vec{x})\right)}_{\text{even}} = 0$$

1. DFs that depend only on \mathcal{H}

Velocity dispersions

$$\sigma_{ij}^2 = \frac{1}{\nu(\bar{x})} \int \underbrace{(v_i - \bar{v})(v_j - \bar{v})}_{=0 \quad =0} f\left(\frac{1}{2} \bar{v}^2 + \phi(\bar{x})\right) d^3\bar{v}$$

odd, except if $i=j$ ($\sigma_{xx} = \sigma_{yy} = \sigma_{zz}$)

$$\sigma^2 = \frac{1}{\nu(\bar{x})} \int_{-\infty}^{\infty} v_z^2 dv_x \int_{-\infty}^{\infty} dv_y \int_{-\infty}^{\infty} dv_z f\left(\frac{1}{2} \bar{v}^2 + \phi(\bar{x})\right)$$

using spherical coord in velocity space :

$$\left\{ \begin{aligned} dv_x dv_y dv_z &= v^2 \sin\theta dv d\theta d\phi \\ v_z^2 &= v^2 \cos^2\theta \\ v^2 &= v_x^2 + v_y^2 + v_z^2 \end{aligned} \right.$$

$$\sigma^2 = \frac{4}{3} \pi \frac{1}{\nu(\bar{x})} \int_0^{\infty} v^4 f\left(\frac{1}{2} v^2 + \phi(\bar{x})\right) dv$$

$$\sigma_{ij} = \begin{pmatrix} \sigma & 0 & 0 \\ 0 & \sigma & 0 \\ 0 & 0 & \sigma \end{pmatrix}$$

isotropic system:
the velocity ellipsoid is a sphere

2. DFs that depend on H and \vec{L}

(spherical symmetry)

$$\phi = \phi(r)$$

We restrict our study to symmetric DFs

: indep. of any direction

$$f(\vec{x}, \vec{v}) = f(H, L)$$



$$\vec{L} \rightarrow |\vec{L}| = L$$

we consider

radial velocity: $\vec{v}_r = v_r \vec{e}_r$

tangential velocity: $\vec{v}_t = \vec{v} - v_r \vec{e}_r$

$$v_t^2 = v_e^2 + v_\varphi^2$$



$$v_e = v_t \cos \theta \quad v_\varphi = v_t \sin \theta$$

$$\left\{ \begin{array}{l} L = r^2 \dot{\theta} = r v_t = r \sqrt{v_e^2 + v_\varphi^2} \\ H = \frac{1}{2} (v_r^2 + v_t^2) + \phi(r) \end{array} \right.$$

2. DFs that depend on H and \vec{L}

Mean velocity

$$\bar{v}_r = \frac{1}{\Psi(r)} \int v_r f\left(\frac{1}{2}(v_r^2 + v_t^2) + \phi(r), r, v_t\right) d^3v$$

$$= \frac{1}{\Psi(r)} \int_{-\infty}^{\infty} \underbrace{v_r}_{\text{odd in } v_r} dv_r \int \underbrace{d^2\vec{v}_t}_{\text{even in } v_r} f\left(\frac{1}{2}(v_r^2 + v_t^2) + \phi(r), r, v_t\right) = 0$$

$$\bar{v}_t = \frac{1}{\Psi(r)} \int \underbrace{\vec{v}_t}_{\text{odd in } v_t} d^2\vec{v}_t \int_{-\infty}^{\infty} \underbrace{dv_r}_{\text{even in } v_t} f\left(\frac{1}{2}(v_r^2 + v_t^2) + \phi(r), r, v_t\right) = 0$$

Note $v_t \in [0, \infty]$

2. DFs that depend on H and \vec{L}

Velocity dispersions

} veloc. in c.g.f. coord
 $dV_e dV_\varphi \rightarrow v_t dV_t$

$$\begin{aligned} \sigma_r^2 &= \frac{1}{V(\infty)} \int_{-\infty}^{\infty} v_r^2 dv_r \int_{-\infty}^{\infty} dV_e \int_{-\infty}^{\infty} dV_\varphi f\left(\frac{1}{2}(v_r^2 + v_e^2 + v_\varphi^2) + \phi(r), r v_t\right) \\ &= \frac{2\pi}{V(\infty)} \int_{-\infty}^{\infty} v_r^2 dv_r \int_0^{\infty} dV_t v_t f\left(\frac{1}{2}(v_r^2 + v_t^2) + \phi(r), r v_t\right) \neq 0 \end{aligned}$$

$$\begin{aligned} \sigma_e^2 &= \frac{1}{V(\infty)} \int_{-\infty}^{\infty} v_e^2 dv_e \int_{-\infty}^{\infty} dV_\varphi \int_{-\infty}^{\infty} dv_r f\left(\frac{1}{2}(v_r^2 + v_e^2 + v_\varphi^2) + \phi(r), r v_t\right) \\ &= \frac{1}{V(\infty)} \int_{-\infty}^{\infty} \int_0^{\infty} v_e^2 v_t dv_t dv_r f\left(\frac{1}{2}(v_r^2 + v_t^2) + \phi(r), r v_t\right) \\ &= \frac{\pi}{V(\infty)} \int_0^{\infty} dv_t v_t^3 \int_{-\infty}^{\infty} dv_r f\left(\frac{1}{2}(v_r^2 + v_t^2) + \phi(r), r v_t\right) \end{aligned}$$

$dV_e dV_\varphi \rightarrow v_t dV_t$

$v_e^2 v_t dv_e = v_t^2 \cos^2 \theta v_t dv_t \rightarrow \pi v_t^3 dv_t$

2. DFs that depend on H and \vec{L}

Velocity dispersions

$$\begin{aligned}
 \sigma_{\varphi}^2 &= \frac{1}{V(x)} \int_{-\infty}^{\infty} v_{\varphi}^2 dv_{\varphi} \int_{-\infty}^{\infty} dv_e \int_{-\infty}^{\infty} dv_r f\left(\frac{1}{2}(v_r^2 + v_e^2 + v_{\varphi}^2) + \phi(r), r v_t\right) \\
 &= \frac{1}{V(x)} \int_{-\infty}^{\infty} \int_0^{\infty} v_{\varphi}^2 v_t dv_t dv_r f\left(\frac{1}{2}(v_r^2 + v_t^2) + \phi(r), r v_t\right) \\
 &= \frac{\pi}{V(x)} \int_0^{\infty} dv_t v_t^3 \int_{-\infty}^{\infty} dv_r f\left(\frac{1}{2}(v_r^2 + v_t^2) + \phi(r), r v_t\right)
 \end{aligned}$$

$dv_e dv_{\varphi} \rightarrow v_t dv_t$
 $v_{\varphi}^2 = v_t^2 \sin^2 \theta \rightarrow \pi v_t^3 dv_t$

$$\sigma_{\varphi}^2 = \sigma_e^2$$



ok, spherical symmetry

$$\sigma_{ij} = 0 \quad \text{if } i \neq j$$

Anisotropic system

$$\sigma_r^2 \neq \sigma_e^2 = \sigma_{\varphi}^2$$

The velocity ellipsoid is

oblate  or prolate 


3. DFs that depend on H and L_z

(cylindrical symmetry)

$$\phi = \phi(R, z)$$

$$f(\vec{x}, \vec{v}) = f(H, L_z)$$

$$\begin{cases} H = \frac{1}{2} (v_r^2 + v_\phi^2 + v_z^2) + \phi(R, z) \\ L_z = R^2 \dot{\phi} = R v_\phi \quad (v_\phi = R \dot{\phi}) \end{cases}$$



Mean velocity

$$\bar{v}_R = \int dv_R v_R \int dv_z dv_\phi f\left(\frac{1}{2} (v_r^2 + v_\phi^2 + v_z^2) + \phi(R, z), R v_\phi\right) = 0$$

odd in v_R

$$\bar{v}_z = \int dv_z v_z \int dv_R dv_\phi f\left(\frac{1}{2} (v_r^2 + v_\phi^2 + v_z^2) + \phi(R, z), R v_\phi\right) = 0$$

odd in v_z

$$\bar{v}_\phi = \int dv_\phi v_\phi \int dv_R dv_z f\left(\frac{1}{2} (v_r^2 + v_\phi^2 + v_z^2) + \phi(R, z), R v_\phi\right)$$

$\neq 0$ in general (net rotation)

$= 0$ only if f is an even function of $L_z = R v_\phi$

3. DFs that depend on H and L_z

Velocity dispersions

$$\sigma_R^2 = \frac{1}{V(\infty)} \int dV_R V_R^2 \int dV_z \int dV_\varphi \rho \left(\frac{1}{2} (V_R^2 + V_\varphi^2 + V_z^2) + \phi(R, z), R V_\varphi \right)$$

$$\sigma_z^2 = \sigma_R^2 \quad (\text{both variables } V_R \text{ and } V_z \text{ can be exchanged})$$

$$\sigma_\varphi^2 = \frac{1}{V(z)} \int dV_\varphi (V_\varphi - \bar{V}_\varphi)^2 \int dV_z dV_R \rho \left(\frac{1}{2} (V_R^2 + V_\varphi^2 + V_z^2) + \phi(R, z), R V_\varphi \right)$$



σ is isotropic in the meridional plane



Anisotropic system

$$\sigma_\varphi^2 \neq \sigma_R^2 = \sigma_z^2$$

The velocity ellipsoid is

oblate  or prolate 

Interpretation

Example 1

1-D potential


$$\left\{ \begin{array}{l} E = \frac{1}{2} v^2 + \phi(r) \\ v = \pm \sqrt{2(E - \phi(r))} \end{array} \right.$$

a) $f(x, v) = f(E) = \delta(E - E_0)$

$$\left\{ \begin{array}{ll} \infty & v = \pm \sqrt{2(E_0 - \phi(r))} \\ 0 & \text{instead} \end{array} \right.$$

b) $f(x, v) = f(E)$

↳
give a weight to
orbits depending on
their energy



Example 2

3D - spherical potential

- orbits described in planes, characterized by (E, L)

a) Ergodic DF :

$$g(\vec{x}, \vec{v}) = g(E)$$

$$\begin{pmatrix} \sigma & 0 & 0 \\ 0 & \sigma & 0 \\ 0 & 0 & \sigma \end{pmatrix}$$

- model built-out of all orbits of all planes with a weight that depends on the energy
(radial and circular orbits are weighted the same way) invariant under rotation (isotropic)

b) non Ergodic DF :

$$g(\vec{x}, \vec{v}) = g(E, L)$$

$$\sigma_r^2 \neq \sigma_\theta^2 = \sigma_\varphi^2$$

- model built-out of all orbits of all planes with a weight that depends on E and L
(radial and circular orbits are weighted differently)

c) non Ergodic DF :

$$g(\vec{x}, \vec{v}) = g(E, \vec{L}) = g_E(E) g_L(\vec{L})$$

$$\text{with } g_L(\vec{L}) = 0 \text{ if } \begin{cases} L_x \neq 0 \\ L_y \neq 0 \end{cases}$$

$$\sigma_\varphi^2 \neq \sigma_r^2 = \sigma_z^2$$

- model built-out of orbits lying in the $z=0$ plane with a weight that depends on E and L_z

The End