

Exercise 9.1. Show that a covector field ξ on a smooth manifold M is smooth if and only if for any smooth vector field X on M the function $\langle \sigma, X \rangle : M \rightarrow \mathbb{R}$ defined by $\langle \sigma, X \rangle(p) = \sigma_p(X_p)$ is smooth.

Exercise 9.2 (Properties of the differential). Let $f, g \in C^\infty(M, \mathbb{R})$.

- Prove the formulas: $d(af + bg) = a df + b dg$ (where a, b are constants), $d(fg) = f dg + g df$, $d(\frac{f}{g}) = \frac{g df - f dg}{g^2}$ (on the set where $g \neq 0$)
- If $h : \mathbb{R} \rightarrow \mathbb{R}$ is a smooth function then $d(h \circ f) = (h' \circ f) df$.
- If $df \equiv 0$, then f is constant on each connected component of M .

Exercise 9.3 (Closed and exact 1-forms). Let M be a smooth manifold, $\xi \in \Omega^1(M)$.

- Show that for every $p \in M$ there exists $f \in C^\infty(M)$ such that $\omega|_p = df|_p$.
Note that this is only an equality of the covectors at one single point p .
- Write $\xi = \sum_i \xi_i d\phi^i$ in some chart (U, ϕ) . Show that if ξ is exact, then

$$\frac{\partial}{\partial \phi^j} \xi_i = \frac{\partial}{\partial \phi^i} \xi_j \quad \text{on } U. \quad (1)$$

- Use the preceding fact to write down a 1-form which is not exact.

*Remark: A 1-form that satisfies (??) for all charts (U, ϕ) is called **closed**. We have just proved that closedness is a necessary condition for exactness. However, it is not always sufficient. The topology of M comes into play: e.g. on a convex subset of \mathbb{R}^n any closed 1-form is exact. But on the punctured plane $\mathbb{R}^2 \setminus \{0\}$ we can construct a closed 1-form that is not exact.*

Exercise 9.4 (A closed 1-form that is not exact). Let $M = \mathbb{R}^2 \setminus \{0\}$. Let $\omega \in \Omega^1(M)$ be given by

$$\omega = \frac{x dy - y dx}{x^2 + y^2}.$$

Compute the integral of ω along the curve

$$\gamma : [0, 2\pi] \rightarrow M : t \mapsto (\cos t, \sin t).$$

Conclude that ω is not exact.

Exercise 9.5. Let (x, y) be the standard coordinates on \mathbb{R}^2 and let (r, φ) be the polar coordinates.

- Express dx and dy in terms of dr and $d\varphi$ (wherever the latter are defined).
- Let $G : \mathbb{R}^2 \rightarrow \mathbb{R}$, $G(x, y) = x^2 + y^2$. Let t be the standard coordinate on \mathbb{R} . Compute $G^*(dt)$.

Exercise 9.6 (Line integrals). .

- Let M be a smooth manifold, $\gamma : I = [a, b] \rightarrow M$ a smooth curve and let $\xi \in \Omega^1(M)$. Denote by t the standard coordinate on \mathbb{R} . Show that $\int_\gamma \xi = \int_I \gamma^* \xi$.
- (Change of variables for 1-forms) Show that if $\sigma : I \rightarrow J$ is a positive (i.e. order preserving) diffeo between two intervals $I = [a, b]$, $J = [c, d]$, then $\int_I \sigma^* \theta = \int_J \theta$ for any 1-form $\theta \in \Omega^1(J)$.

Hint: Compute the derivatives of the functions $F(s) = \int_a^s \sigma^* \theta$ and $G(t) = \int_c^t \theta$.

What happens if σ is a negative (i.e. order reversing) diffeo ?

- (Reparametrization invariance of curve integrals) If two \mathcal{C}^1 curves $\gamma : J \rightarrow M$, $\beta : I \rightarrow M$ are equivalent as oriented curves, in the sense that β is a positive reparametrization of γ (i.e. $\beta = \gamma \circ \sigma$, where $\sigma : I \rightarrow J$ is a positive diffeo), then $\int_\gamma \xi = \int_\beta \xi$ for any 1-form $\xi \in \Omega^1(M)$. Prove this using the definition via pullback.