

**Problem 1**

$B$  is true. If  $w_i(\epsilon) \neq 0$  for all  $i$  then the three arrays have rank  $K$  and there are  $K$  terms in the tensor decomposition. Therefore, by Jennrich's theorem, the decomposition is unique and the rank of the tensor  $T(\epsilon)$  is  $K$ .

$A$  is not true if there exists some  $(i, \epsilon)$  such that  $w_i(\epsilon)$  is zero.

$C$  is not true because all the functions  $w_i$  are continuous. Therefore,  $\lim_{\epsilon \rightarrow 0} T(\epsilon) = T(0)$  and, by Jennrich's theorem, the rank is at most  $K$  (the rank is  $K$  if  $\forall i : w_i(0) \neq 0$ ).

$D$  is not true because if  $w_i(\epsilon) \neq 0$  for all  $i$  and  $\epsilon \in [0, 1]$  then the rank is  $K$ .

**Problem 2**

1. Working with components, we have on one hand

$$T_2^{\alpha\beta} = \sum_{i=1}^k w_i a_i^\alpha a_i^\beta,$$

and on the other hand

$$\begin{aligned} (V \text{Diag}(w_1, \dots, w_k) V^T)^{\alpha\beta} &= \sum_{i,j=1}^n V^{\alpha i} w_i \delta_{ij} (V^T)^{j\beta} = \sum_{i,j=1}^n V^{\alpha i} w_i \delta_{ij} V^{\beta j} \\ &= \sum_{i=1}^n V^{\alpha i} w_i V^{\beta i} = \sum_{i=1}^n \mathbf{a}_i^\alpha w_i \mathbf{a}_i^\beta. \end{aligned}$$

Exactly the same calculation applies to

$$T_3(I, I, \mathbf{u}) = \sum_{i=1}^k w_i (\mathbf{u}^T \mathbf{a}_i) (\mathbf{a}_i \otimes \mathbf{a}_i)$$

with  $w_i$  replaced by  $w_i(\mathbf{u}^T \mathbf{a}_i)$ . It remains to notice that

$$\text{Diag}(w_1(\mathbf{u}^T \mathbf{a}_1), \dots, w_k(\mathbf{u}^T \mathbf{a}_k)) = \text{Diag}(w_1, \dots, w_k) \text{Diag}(\mathbf{u}^T \mathbf{a}_1, \dots, \mathbf{u}^T \mathbf{a}_k).$$

2. When  $n = k$ , since the vectors  $\mathbf{a}_i$  are linearly independent, the matrix  $V$  is square and full rank so invertible. Since the  $w_i$ 's are non-zero, it directly follows that  $T_2$  is also invertible and

$$T_2^{-1} = (V^{-1})^T \text{Diag}\left(\frac{1}{w_1}, \dots, \frac{1}{w_k}\right) V^{-1}.$$

3. a) First, note that:

$$\begin{aligned} M &= T_3(I, I, \mathbf{u})T_2^{-1} \\ &= V\text{Diag}(w_1, \dots, w_k)\text{Diag}(\mathbf{u}^T \mathbf{a}_1, \dots, \mathbf{u}^T \mathbf{a}_k)V^T(V^{-1})^T\text{Diag}\left(\frac{1}{w_1}, \dots, \frac{1}{w_k}\right)V^{-1} \\ &= V\text{Diag}(\mathbf{u}^T \mathbf{a}_1, \dots, \mathbf{u}^T \mathbf{a}_k)V^{-1} \end{aligned}$$

Thus,

$$MV = V\text{Diag}(\mathbf{u}^T \mathbf{a}_1, \dots, \mathbf{u}^T \mathbf{a}_k)$$

which is equivalent to

$$\forall i \in \{1, \dots, K\} : M\mathbf{a}_i = \lambda_i \mathbf{a}_i \quad \text{with} \quad \lambda_i = \mathbf{u}^T \mathbf{a}_i .$$

When  $\mathbf{u}$  is taken at random from a continuous distribution, the inner products  $\mathbf{a}_i^T \mathbf{u}$  are all distinct and nonzero with probability one. Indeed, the set of  $\mathbf{u}$ 's satisfying either  $\mathbf{a}_i^T \mathbf{u} = \mathbf{a}_j^T \mathbf{u}$  for some  $i \neq j$  or  $\mathbf{a}_i^T \mathbf{u} = 0$  for some  $i$  has measure zero. Therefore, we can uniquely determine the (normalized) eigenvectors  $\mathbf{a}_i$ 's by diagonalizing  $M$ .

b) Once we have recovered  $V$ , we can determine the  $w_i$ 's from the diagonal matrix  $V^{-1}T_2(V^{-1})^T$ .

### Problem 3

The first identity simply follows from the definitions:

$$(\mathbf{c} \otimes_{\text{Kro}} \mathbf{b})^T = [c_1 \mathbf{b}^T \quad c_2 \mathbf{b}^T \quad \dots \quad c_J \mathbf{b}^T] = \mathbf{c}^T \otimes_{\text{Kro}} \mathbf{b}^T .$$

For the second identity on the inner product between the two column vectors  $\mathbf{e} \otimes_{\text{Kro}} \mathbf{d}$  and  $\mathbf{c} \otimes_{\text{Kro}} \mathbf{b}$ , we simply have:

$$(\mathbf{e} \otimes_{\text{Kro}} \mathbf{d})^T (\mathbf{c} \otimes_{\text{Kro}} \mathbf{b}) = [e_1 \mathbf{d}^T \quad e_2 \mathbf{d}^T \quad \dots \quad e_J \mathbf{d}^T] \begin{bmatrix} c_1 \mathbf{b} \\ c_2 \mathbf{b} \\ \vdots \\ c_J \mathbf{b} \end{bmatrix} = \sum_{j=1}^J e_j c_j \mathbf{d}^T \mathbf{b} = (\mathbf{e}^T \mathbf{c})(\mathbf{d}^T \mathbf{b}) .$$

Finally, the product of the  $R \times IJ$  matrix  $(E \otimes_{\text{Khr}} D)^T$  and the  $IJ \times R$  matrix  $(C \otimes_{\text{Khr}} B)$  is the  $R \times R$  matrix whose entries are  $\forall (i, j) \in \{1, \dots, R\}^2$ :

$$\begin{aligned} [(E \otimes_{\text{Khr}} D)^T (C \otimes_{\text{Khr}} B)]_{ij} &= \sum_{k=1}^{IJ} [E \otimes_{\text{Khr}} D]_{ki} [C \otimes_{\text{Khr}} B]_{kj} \\ &= (\mathbf{e}_i \otimes_{\text{Kro}} \mathbf{d}_i)(\mathbf{c}_j \otimes_{\text{Kro}} \mathbf{b}_j) \\ &= (\mathbf{e}_i^T \mathbf{c}_j)(\mathbf{d}_i^T \mathbf{b}_j) \\ &= [E^T C]_{ij} [D^T B]_{ij} \\ &= [(E^T C) \circ (D^T B)]_{ij} . \end{aligned}$$

The third equality follows from the identity on the inner product of two Kronecker products. Hence  $(E \otimes_{\text{Khr}} D)^T (C \otimes_{\text{Khr}} B) = (E^T C) \circ (D^T B)$ .