

---

Exercise Set 1  
Quantum Computation

---

**Exercise 1** *Dirac's notation for vectors and matrices*

Let  $\mathcal{H} = \mathbb{C}^N$  be a vector space of  $N$  dimensional vectors with complex components. Let

$$\vec{v} = \begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_N \end{pmatrix}$$

be a column vector. We define its “conjugate” as

$$\vec{v}^\dagger = (\bar{v}_1, \dots, \bar{v}_N)$$

where  $\bar{z}$  stands for the complex conjugate of  $z \in \mathbb{C}$ . So  $\vec{v}^\dagger$  is obtained by transposition and complex conjugation (if the components are real this reduces to the usual transposed vector). The inner or scalar product is by definition

$$\vec{v}^\dagger \cdot \vec{w} = \bar{v}_1 w_1 + \dots + \bar{v}_N w_N$$

and the norm is

$$\|\vec{v}\|^2 = \vec{v}^\dagger \cdot \vec{v} = \bar{v}_1 v_1 + \dots + \bar{v}_N v_N = |v_1|^2 + \dots + |v_N|^2$$

In Dirac's notation we write  $\vec{v} = |v\rangle$  and  $\vec{v}^\dagger = \langle v|$ . Therefore the inner product becomes

$$\langle v|w\rangle = \bar{v}_1 w_1 + \dots + \bar{v}_N w_N$$

The canonical orthonormal basis vectors are by definition

$$\vec{e}_1 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}, \dots, \vec{e}_N = \begin{pmatrix} 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}$$

In Dirac's notation the orthonormality of the basis vectors is expressed as

$$\langle e_i|e_j\rangle = \delta_{ij} = \begin{cases} 1, & \text{if } i = j \\ 0, & \text{if } i \neq j. \end{cases}$$

The expansion of a vector on this basis is

$$|v\rangle = v_1 |e_1\rangle + v_2 |e_2\rangle + \dots + v_N |e_N\rangle$$

Now you will check a few easy facts of linear algebra and translate them in Dirac's notation.

- (a) Check from the definitions above that if  $|v\rangle = \alpha |v'\rangle + \beta |v''\rangle$  then

$$\langle v| = \bar{\alpha} \langle v'| + \bar{\beta} \langle v''|.$$

- (b) In particular deduce that if  $|v\rangle = v_1 |e_1\rangle + v_2 |e_2\rangle + \dots + v_N |e_N\rangle$  then

$$\langle v| = \bar{v}_1 \langle e_1| + \bar{v}_2 \langle e_2| + \dots + \bar{v}_N \langle e_N|.$$

- (c) Show directly in Dirac notation that

$$\langle v|w\rangle = \bar{v}_1 w_1 + \dots + \bar{v}_N w_N.$$

- (d) Deduce that  $\sqrt{\langle v|v\rangle} = \|v\|$ .

- (e) Consider the ket-bra expression  $|e_i\rangle \langle e_j|$  for canonical basis vectors. Write this as an  $N \times N$  matrix.

- (f) Consider now an  $N \times N$  matrix  $A$  with complex matrix elements  $a_{ij}$ ;  $i = 1 \dots N$ ;  $j = 1 \dots N$ . Deduce from the above question that

$$A = \sum_{i,j=1}^N a_{ij} |e_i\rangle \langle e_j|$$

and also that

$$a_{ij} = \langle e_i| A |e_j\rangle.$$

- (g) In particular verify that the identity matrix satisfies :

$$I = \sum_{i=1}^N |e_i\rangle \langle e_i|.$$

This is called the closure relation. Deduce that in fact this relation is valid for any orthonormal basis of vectors  $|\varphi_i\rangle$ ,  $i = 1, \dots, N$ .

- (h) (Spectral theorem) Let  $A = A^\dagger$  where  $A^\dagger = A^{T,*}$ . This is called a *Hermitian matrix*. An important theorem of linear algebra states that : “any Hermitian matrix has  $N$  orthonormal eigenvectors with real eigenvalues (possibly degenerate)”. Let  $|\varphi_i\rangle$ ,  $\alpha_i$ ,  $i = 1, \dots, N$  be the eigenvectors and eigenvalues of  $A$ , *i.e.*,

$$A |\varphi_i\rangle = \alpha_i |\varphi_i\rangle.$$

Prove directly in Dirac’s notation that

$$A = \sum_{i=1}^N \alpha_i |\varphi_i\rangle \langle \varphi_i|.$$

This “expansion” is often called the spectral expansion (or theorem).

**Exercise 2** *Tensor Product in Dirac's notation*

Let  $\mathcal{H}_1 = \mathbb{C}^N$  and  $\mathcal{H}_2 = \mathbb{C}^M$  be  $N$  and  $M$  dimensional Hilbert spaces. The tensor product space  $\mathcal{H}_1 \otimes \mathcal{H}_2$  is a new Hilbert space formed by "pairs of vectors" denoted as  $|v\rangle_1 \otimes |w\rangle_2 \equiv |v, w\rangle$  with the properties :

- $(\alpha |v\rangle_1 + \beta |v'\rangle_1) \otimes |w\rangle_2 = \alpha |v\rangle_1 \otimes |w\rangle_2 + \beta |v'\rangle_1 \otimes |w\rangle_2,$
- $|v\rangle_1 \otimes (\alpha |w\rangle_2 + \beta |w'\rangle_2) = \alpha |v\rangle_1 \otimes |w\rangle_2 + \beta |v\rangle_1 \otimes |w'\rangle_2,$
- $(|v\rangle_1 \otimes |w\rangle_2)^\dagger = \langle v|_1 \otimes \langle w|_2,$
- $\langle v, w | v', w' \rangle = \langle v | v' \rangle_1 \langle w | w' \rangle_2.$

- (a) Show that for any two vectors of  $\mathcal{H}_1$  and  $\mathcal{H}_2$  expanded on two basis,  $|v\rangle_1 = \sum_{i=1}^N v_i |e_i\rangle_1$  and  $|w\rangle_2 = \sum_{j=1}^M w_j |f_j\rangle_2$ , then

$$|v\rangle_1 \otimes |w\rangle_2 = \sum_{i=1}^N \sum_{j=1}^M v_i w_j |e_i\rangle_1 \otimes |f_j\rangle_2.$$

- (b) Show that if  $\{|e_i\rangle_1; i = 1 \dots N\}$  and  $\{|f_j\rangle_2; j = 1 \dots M\}$  are orthonormal, then  $|e_i\rangle_1 \otimes |f_j\rangle_2 \equiv |e_i, f_j\rangle$  is an orthonormal basis of  $\mathcal{H}_1 \otimes \mathcal{H}_2$ . What is the dimension of  $\mathcal{H}_1 \otimes \mathcal{H}_2$ ?
- (c) Any vector  $|\Psi\rangle$  of  $\mathcal{H}_1 \otimes \mathcal{H}_2$  can be expanded on the basis  $|e_i\rangle_1 \otimes |f_j\rangle_2 \equiv |e_i, f_j\rangle, i = 1 \dots N, j = 1 \dots M,$

$$|\Psi\rangle = \sum_{i=1, j=1}^{N, M} \psi_{ij} |e_i, f_j\rangle.$$

If  $A$  is a matrix acting on  $\mathcal{H}_1$  and  $B$  is a matrix acting on  $\mathcal{H}_2$ , the tensor product  $A \otimes B$  is defined as

$$A \otimes B |\Psi\rangle = \sum_{i, j} \psi_{ij} A |e_i\rangle_1 \otimes B |f_j\rangle_2.$$

Check that the matrix elements of  $A \otimes B$  in the basis  $|e_i, f_j\rangle$  are :

$$\langle e_i, f_j | A \otimes B | e_k, f_l \rangle = a_{ik} b_{jl}.$$

- (d) Let  $\mathcal{H}_1 = \mathbb{C}^2, \mathcal{H}_2 = \mathbb{C}^2$ . Take  $A_1 = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, B_2 = \begin{pmatrix} e & f \\ g & h \end{pmatrix}, |v\rangle_1 = \begin{pmatrix} \alpha \\ \beta \end{pmatrix}, |w\rangle_2 = \begin{pmatrix} \gamma \\ \delta \end{pmatrix}.$   
From the defining properties of the tensor product deduce the the following formulas :

$$|v\rangle_1 \otimes |w\rangle_2 = \begin{pmatrix} \alpha\gamma \\ \alpha\delta \\ \beta\gamma \\ \beta\delta \end{pmatrix}, \quad A_1 \otimes B_2 = \begin{pmatrix} ae & af & be & bf \\ ag & ah & bg & bh \\ ce & cf & de & df \\ cg & ch & dg & dh \end{pmatrix}.$$

These are often useful in order to do calculations in components.