

---

Homework 9  
CS-526 Learning Theory

---

**Note:** The tensor product is denoted by  $\otimes$ . In other words, for vectors  $\mathbf{a}, \mathbf{b}, \mathbf{c}$  we have that  $\mathbf{a} \otimes \mathbf{b}$  is the square array  $a^\alpha b^\beta$  where the superscript denotes the components, and  $\mathbf{a} \otimes \mathbf{b} \otimes \mathbf{c}$  is the cubic array  $a^\alpha b^\beta c^\gamma$ . We often denote components by superscripts because we need the lower index to label vectors themselves.

**Problem 1: A multiple choice question**

Find the correct answer(s).

Let  $w_i(\epsilon)$  for  $i \in \{1, \dots, K\}$  be continuous functions of  $\epsilon \in [0, 1]$ . Suppose that for all  $\epsilon \in [0, 1]$  the  $N \times K$  matrices  $[\mathbf{a}_1 + \epsilon \mathbf{a}'_1 \ \cdots \ \mathbf{a}_K + \epsilon \mathbf{a}'_K]$ ,  $[\mathbf{b}_1 + \epsilon \mathbf{b}'_1 \ \cdots \ \mathbf{b}_K + \epsilon \mathbf{b}'_K]$  and  $[\mathbf{c}_1 + \epsilon \mathbf{c}'_1 \ \cdots \ \mathbf{c}_K + \epsilon \mathbf{c}'_K]$  have rank  $K$ . Consider the tensor

$$T(\epsilon) = \sum_{i=1}^K w_i(\epsilon) (\mathbf{a}_i + \epsilon \mathbf{a}'_i) \otimes (\mathbf{b}_i + \epsilon \mathbf{b}'_i) \otimes (\mathbf{c}_i + \epsilon \mathbf{c}'_i) .$$

- (A) The tensor rank equals  $K$  for all  $\epsilon \in [0, 1]$ .
- (B) The tensor rank equals  $K$  for all  $\epsilon \in [0, 1]$  such that  $\forall i \in \{1, \dots, K\} : w_i(\epsilon) \neq 0$ .
- (C) It may happen that the tensor rank of the limit  $\lim_{\epsilon \rightarrow 0} T(\epsilon)$  is  $K + 1$ .
- (D) If we replace the assumption that  $[\mathbf{c}_1 + \epsilon \mathbf{c}'_1 \ \cdots \ \mathbf{c}_K + \epsilon \mathbf{c}'_K]$  is rank  $K$  by the assumption that these vectors are pairwise independent, then the tensor rank can *never* be  $K$  whatever the assumptions on  $w_i(\epsilon)$ ,  $i = 1, \dots, K$ .

**Problem 2: Kronecker, Khatri-Rao, Hadamard products: check useful identities**

We recall a few definitions seen in class. The Kronecker product of two column vectors  $\mathbf{b} \in \mathbb{R}^I$  and  $\mathbf{c} \in \mathbb{R}^J$  is the column vector:

$$\mathbf{c} \otimes_{\text{Kro}} \mathbf{b} \triangleq [c_1 \mathbf{b}^T \ c_2 \mathbf{b}^T \ \cdots \ c_J \mathbf{b}^T]^T .$$

The Kronecker product of two row vectors  $\mathbf{d}$  and  $\mathbf{e}$  is the row vector:

$$\mathbf{d} \otimes_{\text{Kro}} \mathbf{e} \triangleq [d_1 \mathbf{e} \ d_2 \mathbf{e} \ \cdots \ d_J \mathbf{e}] .$$

The Khatri-Rao product of two matrices  $B = [\mathbf{b}_1 \ \mathbf{b}_2 \ \cdots \ \mathbf{b}_R] \in \mathbb{R}^{I \times R}$  and  $C = [\mathbf{c}_1 \ \mathbf{c}_2 \ \cdots \ \mathbf{c}_R] \in \mathbb{R}^{J \times R}$  is the  $(IJ) \times R$  matrix:

$$C \otimes_{\text{Khr}} B \triangleq [\mathbf{c}_1 \otimes_{\text{Kro}} \mathbf{b}_1 \ \cdots \ \mathbf{c}_R \otimes_{\text{Kro}} \mathbf{b}_R] .$$

Finally, the Hadamard product of two matrices (of same dimensions) is the matrix given by the point-wise product of components, i.e, if  $A, B$  have matrix elements  $a_{ij}$  and  $b_{ij}$  then the Hadamard product  $A \circ B$  has matrix elements  $a_{ij}b_{ij}$ .

Let  $\mathbf{b}, \mathbf{d} \in \mathbb{R}^I$  and  $\mathbf{c}, \mathbf{e} \in \mathbb{R}^J$  be column vectors. Let  $B, D \in \mathbb{R}^{I \times R}$  and  $C, E \in \mathbb{R}^{J \times R}$  be four matrices. Check the following identities used in class:

$$\begin{aligned}(\mathbf{c} \otimes_{\text{Kro}} \mathbf{b})^T &= \mathbf{c}^T \otimes_{\text{Kro}} \mathbf{b}^T ; \\(\mathbf{e} \otimes_{\text{Kro}} \mathbf{d})^T (\mathbf{c} \otimes_{\text{Kro}} \mathbf{b}) &= (\mathbf{e}^T \mathbf{c})(\mathbf{d}^T \mathbf{b}) ; \\(E \otimes_{\text{Khr}} D)^T (C \otimes_{\text{Khr}} B) &= (E^T C) \circ (D^T B) .\end{aligned}$$